

ON ASYMPTOTIC BEHAVIOR OF SOLUTIONS TO SCHRÖDINGER
EQUATIONS WITH SINGULAR DIPOLE–TYPE POTENTIALS

VERONICA FELLI – ELSA M. MARCHINI – SUSANNA TERRACINI

QUADERNO N. 2/2007 (arxiv:math/0701292v1)



STAMPATO NEL MESE DI APRILE 2007
PRESSO IL DIPARTIMENTO DI MATEMATICA E APPLICAZIONI,
UNIVERSITÀ DEGLI STUDI DI MILANO-BICOCCA, VIA R. COZZI 53, 20125 MILANO, ITALIA.

DISPONIBILE IN FORMATO ELETTRONICO SUL SITO www.matapp.unimib.it.
SEGRETERIA DI REDAZIONE: Ada Osmetti - Giuseppina Cogliandro
tel.: +39 02 6448 5755-5758 fax: +39 02 6448 5705

**Esemplare fuori commercio per il deposito legale agli effetti della Legge 15 aprile 2004
n.106.**

ON ASYMPTOTIC BEHAVIOR OF SOLUTIONS TO SCHRÖDINGER EQUATIONS WITH SINGULAR DIPOLE-TYPE POTENTIALS

VERONICA FELLI, ELSA M. MARCHINI, AND SUSANNA TERRACINI

ABSTRACT. Asymptotics of solutions to Schrödinger equations with singular dipole-type potentials is investigated. We evaluate the exact behavior near the singularity of solutions to elliptic equations with potentials which are purely angular multiples of radial inverse-square functions. Both the linear and the semilinear (critical and subcritical) cases are considered.

Dedicated to Prof. Norman Dancer on the occasion of his 60th birthday.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

In nonrelativistic molecular physics, the interaction between an electric charge and the dipole moment $\mathbf{D} \in \mathbb{R}^N$ of a molecule is described by an inverse square potential with an anisotropic coupling strength. In particular the Schrödinger equation for the wave function of an electron interacting with a polar molecule (supposed to be point-like) can be written as

$$\left(-\frac{\hbar^2}{2m} \Delta + e \frac{x \cdot \mathbf{D}}{|x|^3} - E \right) \Psi = 0,$$

where e and m denote respectively the charge and the mass of the electron and \mathbf{D} is the dipole moment of the molecule, see [11].

We aim to describe the asymptotic behavior near the singularity of solutions to equations associated to dipole-type Schrödinger operators of the form

$$L_{\lambda, \mathbf{d}} := -\Delta - \frac{\lambda(x \cdot \mathbf{d})}{|x|^3}$$

in \mathbb{R}^N , where $N \geq 3$, $\lambda = \frac{2m\epsilon|\mathbf{D}|}{\hbar^2}$, being $|\mathbf{D}|$ the magnitude of the dipole moment \mathbf{D} , and $\mathbf{d} = \mathbf{D}/|\mathbf{D}|$ denotes the orientation of \mathbf{D} . A precise estimate of such a behavior is the first step in the analysis of fundamental properties of Schrödinger operators, such as positivity, essential self-adjointness, and spectral features. Such an analysis has been carried out in [7] for the case of Schrödinger operators with multipolar Hardy potentials based on the regularity results proved in [8] for degenerate elliptic equations. In a forthcoming paper, the authors will apply Theorem 1.1 to establish spectral properties of multi-singular dipole type operators.

We emphasize that, from the mathematical point of view, potentials of the form $\frac{\lambda(x \cdot \mathbf{d})}{|x|^3}$ have the same order of homogeneity as inverse square potentials and consequently share many features

First and third author supported by Italy MIUR, national project “Variational Methods and Nonlinear Differential Equations”.

2000 *Mathematics Subject Classification.* 35J10, 35B40, 35J60.

Keywords. Singular potentials, dipole moment, Hardy’s inequality, Schrödinger operators.

with them, such as invariance by scaling and Kelvin transform, as well as no inclusion in the Kato class. We mention that Schrödinger equations with Hardy-type singular potentials have been largely studied, see e.g. [1, 6, 9, 10, 17, 18] and references therein.

More precisely, in this paper we deal with a more general class of Schrödinger operators including those with dipole-type potentials, namely with operators whose potentials are purely angular multiples of radial inverse-square potentials:

$$\mathcal{L}_a := -\Delta - \frac{a(x/|x|)}{|x|^2}$$

in \mathbb{R}^N , where $N \geq 3$ and $a \in L^\infty(\mathbb{S}^{N-1})$.

The problem of establishing the asymptotic behavior of solutions to elliptic equations near an isolated singular point has been studied by several authors in a variety of contexts, see e.g. [13] for Fuchsian type elliptic operators and [5] for Fuchsian type weighted operators. The asymptotics we derive in this work is not contained in the aforementioned papers, which prove the existence of the limit at the singularity of any quotient of two positive solutions in some linear and semilinear cases which, however, do not include the perturbed linear case and the critical nonlinear case treated here. Moreover, besides proving the existence of such a limit, we also obtain a Cauchy type representation formula for it, see (4) and (8).

We also quote [12], where asymptotics at infinity is established for perturbed inverse square potentials and in some particular nonradial case. Hölder continuity results for degenerate elliptic equations with singular weights and asymptotic analysis of the behavior of solutions near the pole are contained in [8].

As a natural setting to study the properties of operators \mathcal{L}_a , we introduce the functional space $\mathcal{D}^{1,2}(\mathbb{R}^N)$ defined as the completion of $C_c^\infty(\mathbb{R}^N)$ with respect to the Dirichlet norm

$$\|u\|_{\mathcal{D}^{1,2}(\mathbb{R}^N)} := \left(\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx \right)^{1/2}.$$

In order to discuss the positivity properties of the Schrödinger operator \mathcal{L}_a in $\mathcal{D}^{1,2}(\mathbb{R}^N)$, we consider the best constant in the associated Hardy-type inequality

$$(1) \quad \Lambda_N(a) := \sup_{u \in \mathcal{D}^{1,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |x|^{-2} a(x/|x|) u^2(x) dx}{\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx}.$$

By the classical Hardy inequality, $\Lambda_N(a) \leq \frac{4}{(N-2)^2} \operatorname{ess\,sup}_{\mathbb{S}^{N-1}} a$, where $\operatorname{ess\,sup}_{\mathbb{S}^{N-1}} a$ denotes the essential supremum of a in \mathbb{S}^{N-1} . We also notice that, in the dipole case, by rotation invariance, $\Lambda_N\left(\frac{\lambda x}{|x|} \cdot \mathbf{d}\right)$ does not depend on \mathbf{d} .

It is easy to verify that the quadratic form associated to \mathcal{L}_a is positive definite in $\mathcal{D}^{1,2}(\mathbb{R}^N)$ if and only if $\Lambda_N(a) < 1$. The relation between the value $\Lambda_N(a)$ and the first eigenvalue of the angular component of the operator on the sphere \mathbb{S}^{N-1} is discussed in section 2, see Lemmas 2.4 and 2.5. More precisely, Lemma 2.5 ensures that the quadratic form associated to \mathcal{L}_a is positive definite if and only if

$$\mu_1 > -\left(\frac{N-2}{2}\right)^2,$$

where $\mu_1 = \mu_1(a, N)$ is the first eigenvalue of the operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ on \mathbb{S}^{N-1} (see Lemma 2.1). We denote by ψ_1 the associated positive L^2 -normalized eigenfunction and set

$$(2) \quad \sigma = \sigma(a, N) := -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1}.$$

In the spirit of the well-known Riemann removable singularity theorem, we describe the behavior of solutions to linear Schrödinger equations with a dipole-type singularity localized in a neighborhood of 0.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^N$ be a bounded open set such that $0 \in \Omega$. Assume that $q \in L_{\text{loc}}^\infty(\Omega \setminus \{0\})$ is such that $q(x) = O(|x|^{-(2-\varepsilon)})$ as $|x| \rightarrow 0$ for some $\varepsilon > 0$, $a \in L^\infty(\mathbb{S}^{N-1})$ satisfies $\Lambda_N(a) < 1$, and $u \in H^1(\Omega)$, $u \geq 0$ a.e. in Ω , $u \not\equiv 0$, weakly solves*

$$(3) \quad -\Delta u(x) - \frac{a(x/|x|)}{|x|^2} u(x) = q(x) u(x) \quad \text{in } \Omega,$$

i.e.

$$\int_{\Omega} \nabla u(x) \cdot \nabla w(x) dx - \int_{\Omega} \frac{a(x/|x|)}{|x|^2} u(x) w(x) dx = \int_{\Omega} q(x) u(x) w(x) dx,$$

for all $w \in H_0^1(\Omega)$. Then the function

$$x \mapsto \frac{u(x)}{|x|^\sigma \psi_1(x/|x|)}$$

is continuous in Ω and

$$(4) \quad \lim_{|x| \rightarrow 0} \frac{u(x)}{|x|^\sigma \psi_1\left(\frac{x}{|x|}\right)} = \int_{\mathbb{S}^{N-1}} \left(R^{-\sigma} u(R\theta) + \int_0^R \frac{s^{1-\sigma}}{2\sigma + N - 2} q(s\theta) u(s\theta) ds \right. \\ \left. - R^{-2\sigma - N + 2} \int_0^R \frac{s^{N-1+\sigma}}{2\sigma + N - 2} q(s\theta) u(s\theta) ds \right) \psi_1(\theta) dV(\theta),$$

for all $R > 0$ such that $\overline{B(0, R)} := \{x \in \mathbb{R}^N : |x| \leq R\} \subset \Omega$.

We notice that (4) is actually a *Cauchy's integral type formula* for u . Moreover the term at the right hand side is independent of R . In the case in which the perturbation q is radial then an analogous formula holds also for changing sign solutions to (3), see Remark 2.

If the perturbing potential q satisfies some proper summability condition, instead of the stronger control on the blow-up rate at the singularity required in Theorem 1.1, a Brezis-Kato type argument, see [2], allows us to derive an upper estimate on the behavior of solutions. For any $q \geq 1$, we denote as $L^q(\varphi^{2^*}, \Omega)$ the weighted L^q -space endowed with the norm

$$\|u\|_{L^q(\varphi^{2^*}, \Omega)} := \left(\int_{\Omega} \varphi^{2^*}(x) |u(x)|^q dx \right)^{1/q},$$

where $2^* = \frac{2N}{N-2}$ is the critical Sobolev exponent and φ denotes the weight function

$$(5) \quad \varphi(x) := |x|^\sigma \psi_1(x/|x|).$$

The following Brezis-Kato type result holds.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^N$ be a bounded open set such that $0 \in \Omega$, $a \in L^\infty(\mathbb{S}^{N-1})$ satisfying $\Lambda_N(a) < 1$, and $V \in L^s(\varphi^{2^*}, \Omega)$ for some $s > N/2$. Then, for any $\Omega' \Subset \Omega$, there exists a positive constant*

$$C = C(N, a, \|V\|_{L^s(\varphi^{2^*}, \Omega)}, \text{dist}(\Omega', \partial\Omega), \text{diam } \Omega)$$

depending only on N , a , $\|V\|_{L^s(\varphi^{2^}, \Omega)}$, $\text{dist}(\Omega', \partial\Omega)$, and $\text{diam } \Omega$, such that for any weak $H^1(\Omega)$ -solution u of*

$$(6) \quad -\Delta u(x) - \frac{a(x/|x|)}{|x|^2} u(x) = \varphi^{2^*-2}(x)V(x)u(x), \quad \text{in } \Omega,$$

i.e. for any $u \in H^1(\Omega)$ satisfying

$$\int_{\Omega} \nabla u(x) \cdot \nabla w(x) dx - \int_{\Omega} \frac{a(x/|x|)}{|x|^2} u(x)w(x) dx = \int_{\Omega} \varphi^{2^*-2}(x)V(x)u(x)w(x) dx,$$

for all $w \in H_0^1(\Omega)$, there holds $\frac{u}{\varphi} \in L^\infty(\Omega')$ and

$$\left\| \frac{u}{\varphi} \right\|_{L^\infty(\Omega')} \leq C \|u\|_{L^{2^*}(\Omega)}.$$

The Brezis-Kato procedure can be applied also to semilinear problems with at most critical growth, thus providing an upper bound for solutions and then reducing the semilinear problem to a linear one with enough control on the potential at the singularity to apply Theorem 1.1 and to recover the exact asymptotic behavior.

Theorem 1.3. *Let Ω be a bounded open set containing 0 , $a \in L^\infty(\mathbb{S}^{N-1})$ satisfying $\Lambda_N(a) < 1$, and $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ such that, for some positive constant C ,*

$$\left| \frac{f(x, u)}{u} \right| \leq C (1 + |u|^{2^*-2}) \quad \text{for a.e. } (x, u) \in \Omega \times \mathbb{R}.$$

Assume that $u \in H^1(\Omega)$, $u \geq 0$ a.e. in Ω , $u \not\equiv 0$, weakly solves

$$(7) \quad -\Delta u(x) - \frac{a(x/|x|)}{|x|^2} u(x) = f(x, u(x)), \quad \text{in } \Omega,$$

i.e.

$$\int_{\Omega} \nabla u(x) \cdot \nabla w(x) dx - \int_{\Omega} \frac{a(x/|x|)}{|x|^2} u(x)w(x) dx = \int_{\Omega} f(x, u(x))w(x) dx,$$

for all $w \in H_0^1(\Omega)$. Then the function

$$x \mapsto \frac{u(x)}{|x|^\sigma \psi_1(x/|x|)}$$

is continuous in Ω and

$$(8) \quad \lim_{|x| \rightarrow 0} \frac{u(x)}{|x|^\sigma \psi_1\left(\frac{x}{|x|}\right)} = \int_{\mathbb{S}^{N-1}} \left(r^{-\sigma} u(r\theta) + \int_0^r \frac{s^{1-\sigma}}{2\sigma + N - 2} f(s\theta, u(s\theta)) ds \right. \\ \left. - r^{-2\sigma - N + 2} \int_0^r \frac{s^{N-1+\sigma}}{2\sigma + N - 2} f(s\theta, u(s\theta)) ds \right) \psi_1(\theta) dV(\theta),$$

for all $r > 0$ such that $\overline{B(0, r)} := \{x \in \mathbb{R}^N : |x| \leq r\} \subset \Omega$.

Notation. We list below some notation used throughout the paper.

- $B(a, r)$ denotes the ball $\{x \in \mathbb{R}^N : |x - a| < r\}$ in \mathbb{R}^N with center at a and radius r .
- dV denotes the volume element on the sphere \mathbb{S}^{N-1} .
- ω_N denotes the volume of the unit sphere \mathbb{S}^{N-1} , i.e. $\omega_N = \int_{\mathbb{S}^{N-1}} dV(\theta)$.
- for any $a \in L^1(\mathbb{S}^{N-1})$, $\mathbf{f}_{\mathbb{S}^{N-1}} a(\theta) dV(\theta)$ denotes the mean of a on \mathbb{S}^{N-1} , i.e.

$$\mathbf{f}_{\mathbb{S}^{N-1}} a(\theta) dV(\theta) = \frac{1}{\omega_N} \int_{\mathbb{S}^{N-1}} a(\theta) dV(\theta).$$

- the symbol ess sup stands for essential supremum.

2. SPECTRUM OF THE ANGULAR COMPONENT

Due to the structure of the dipole-type potential of equation (3), a natural approach to describe the solutions seems to be the separation of variables. To employ such a technique, we need, as a starting point, the description of the spectrum of the angular part of dipole Schrödinger operators.

Lemma 2.1. *Let $a \in L^\infty(\mathbb{S}^{N-1})$. Then the operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ on \mathbb{S}^{N-1} admits a diverging sequence of eigenvalues $\mu_1 < \mu_2 \leq \dots \leq \mu_k < \dots$ the first of which has the following properties:*

- (i) μ_1 is simple;
- (ii) μ_1 can be characterized as

$$(9) \quad \mu_1 = \min_{\psi \in H^1(\mathbb{S}^{N-1}) \setminus \{0\}} \frac{\int_{\mathbb{S}^{N-1}} |\nabla_{\mathbb{S}^{N-1}} \psi(\theta)|^2 dV(\theta) - \int_{\mathbb{S}^{N-1}} a(\theta) \psi^2(\theta) dV(\theta)}{\int_{\mathbb{S}^{N-1}} \psi^2(\theta) dV(\theta)};$$

- (iii) μ_1 is attained by a C^1 positive eigenfunction ψ_1 such that $\min_{\mathbb{S}^{N-1}} \psi_1 > 0$;
- (iv) if, for some $\kappa \in \mathbb{R}$, $a(\theta) = \kappa$ for a.e. $\theta \in \mathbb{S}^{N-1}$, then $\mu_1 = -\kappa$.
- (v) if a is not constant, then $-\text{ess sup}_{\mathbb{S}^{N-1}} a < \mu_1 < -\mathbf{f}_{\mathbb{S}^{N-1}} a(\theta) dV(\theta)$.

Proof. We prove assertion (v), being (i), (ii), (iii), and (iv) quite standard.

Since the function $\psi \equiv 1$ satisfies

$$\int_{\mathbb{S}^{N-1}} |\nabla_{\mathbb{S}^{N-1}} \psi(\theta)|^2 dV(\theta) - \int_{\mathbb{S}^{N-1}} a(\theta) \psi^2(\theta) dV(\theta) = - \int_{\mathbb{S}^{N-1}} a(\theta) dV(\theta),$$

we deduce that $\mu_1 \leq -\mathbf{f}_{\mathbb{S}^{N-1}} a(\theta) dV(\theta)$. In order to prove the strict inequality, we argue by contradiction and assume that $\mu_1 = -\mathbf{f}_{\mathbb{S}^{N-1}} a(\theta) dV(\theta)$. Then $\psi_1 \equiv 1$ attains the minimum value in (9) but, since a is not constant, it does not satisfy equation $-\Delta_{\mathbb{S}^{N-1}} \psi_1 - a(\theta) \psi_1 = 0$, a contradiction. We can thereby conclude that $\mu_1 < -\mathbf{f}_{\mathbb{S}^{N-1}} a(\theta) dV(\theta)$.

From (9), [18, Lemma 1.1], and the optimality of the best constant in Hardy's inequality, it follows that

$$\begin{aligned} \mu_1 &> \inf_{\psi \in H^1(\mathbb{S}^{N-1}) \setminus \{0\}} \frac{\int_{\mathbb{S}^{N-1}} |\nabla_{\mathbb{S}^{N-1}} \psi(\theta)|^2 dV(\theta) - \operatorname{ess\,sup}_{\mathbb{S}^{N-1}} a \int_{\mathbb{S}^{N-1}} \psi^2(\theta) dV(\theta)}{\int_{\mathbb{S}^{N-1}} \psi^2(\theta) dV(\theta)} \\ &= \inf_{u \in \mathcal{D}^{1,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx - \left(\left(\frac{N-2}{2} \right)^2 + \operatorname{ess\,sup}_{\mathbb{S}^{N-1}} a \right) \int_{\mathbb{R}^N} \frac{u^2(x)}{|x|^2} dx}{\int_{\mathbb{R}^N} \frac{u^2(x)}{|x|^2} dx} \\ &= - \operatorname{ess\,sup}_{\mathbb{S}^{N-1}} a, \end{aligned}$$

thus proving the left part of the inequality stated in (v). \square

The asymptotic behavior of eigenvalues μ_k as $k \rightarrow +\infty$ is described by Weyl's law, which is recalled in the theorem below. We refer to [14, 15] for a proof.

Theorem 2.2 (Weyl's law). *For $a \in L^\infty(\mathbb{S}^{N-1})$, let $\{\mu_k\}_{k \geq 1}$ be the eigenvalues of the operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ on \mathbb{S}^{N-1} . Then*

$$(10) \quad \mu_k = C(N, a) k^{2/(N-1)} (1 + o(1)) \quad \text{as } k \rightarrow +\infty,$$

for some positive constant $C(N, a)$ depending only on N and a .

The following lemma provides an estimate of the L^∞ -norm of eigenfunctions of the operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ in terms of the corresponding eigenvalues. For classical results about L^∞ estimates of eigenfunctions of Schrödinger operators we refer to [16, §4] and references therein.

Lemma 2.3. *For $a \in L^\infty(\mathbb{S}^{N-1})$ and $k \in \mathbb{N} \setminus \{0\}$, let ψ_k be a L^2 -normalized eigenfunction of the Schrödinger operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ on the sphere associated to the k -th eigenvalue μ_k , i.e.*

$$(11) \quad \begin{cases} -\Delta_{\mathbb{S}^{N-1}} \psi_k(\theta) - a(\theta) \psi_k(\theta) = \mu_k \psi_k(\theta), & \text{in } \mathbb{S}^{N-1}, \\ \int_{\mathbb{S}^{N-1}} |\psi_k(\theta)|^2 dV(\theta) = 1. \end{cases}$$

Then, there exists a constant C_1 depending only on N and a such that

$$|\psi_k(\theta)| \leq C_1 |\mu_k|^{\lfloor (N-1)/4 \rfloor + 1},$$

where $\lfloor \cdot \rfloor$ denotes the floor function, i.e. $\lfloor x \rfloor := \min\{j \in \mathbb{Z} : j \leq x\}$.

Proof. Using classical elliptic regularity theory and bootstrap methods, we can easily prove that for any $j \in \mathbb{N}$ there exists a constant $C(N, j)$, depending only on j and N , such that

$$\|\psi_k\|_{W^{2, \frac{2(N-1)}{(N-1)-4(j-1)}}(\mathbb{S}^{N-1})} \leq C(N, j) (\mu_k + \|a\|_{L^\infty(\mathbb{S}^{N-1})})^j.$$

Choosing $j = \lfloor \frac{N-1}{4} \rfloor + 1$, by Sobolev's inclusions we deduce that

$$W^{2, \frac{2(N-1)}{(N-1)-4(j-1)}}(\mathbb{S}^{N-1}) \hookrightarrow C^{0, \alpha}(\mathbb{S}^{N-1}) \hookrightarrow L^\infty(\mathbb{S}^{N-1}),$$

for any $0 < \alpha < 2(1 - \frac{N-1}{4} + \lfloor \frac{N-1}{4} \rfloor)$, thus implying the required estimate. \square

Arguing as in the proof of [18, Lemma 1.1], we can deduce the following characterization of $\Lambda_N(a)$.

Lemma 2.4. For $a \in L^\infty(\mathbb{S}^{N-1})$, let $\Lambda_N(a)$ be defined in (1). Then

$$(12) \quad \Lambda_N(a) = \max_{\substack{\psi \in H^1(\mathbb{S}^{N-1}) \\ \psi \neq 0}} \frac{\int_{\mathbb{S}^{N-1}} a(\theta) \psi^2(\theta) dV(\theta)}{\int_{\mathbb{S}^{N-1}} |\nabla_{\mathbb{S}^{N-1}} \psi(\theta)|^2 dV(\theta) + \left(\frac{N-2}{2}\right)^2 \int_{\mathbb{S}^{N-1}} \psi^2(\theta) dV(\theta)}.$$

We notice that the supremum in (12) is achieved due to the compactness of the embedding $H^1(\mathbb{S}^{N-1}) \hookrightarrow L^2(\mathbb{S}^{N-1})$. As a direct consequence of the above lemma, it is possible to compare $\Lambda_N(a)$ with the best constant in Hardy's inequality

$$\frac{4}{(N-2)^2} = \sup_{u \in \mathcal{D}^{1,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} u^2(x) |x|^{-2} dx}{\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx}.$$

Indeed, if a is not constant, there holds

$$\frac{4}{(N-2)^2} \int_{\mathbb{S}^{N-1}} a(\theta) dV(\theta) < \Lambda_N(a) < \operatorname{ess\,sup}_{\mathbb{S}^{N-1}} a \frac{4}{(N-2)^2},$$

whereas, if $a(\theta) = \kappa$ for a.e. $\theta \in \mathbb{S}^{N-1}$ and for some $\kappa \in \mathbb{R}$, then $\Lambda_N(a) = \frac{4\kappa}{(N-2)^2}$.

Let us consider the quadratic form associated to the Schrödinger operator \mathcal{L}_a , i.e.

$$Q_a(u) := \int_{\mathbb{R}^N} |\nabla u(x)|^2 dx - \int_{\mathbb{R}^N} \frac{a(x/|x|) u^2(x)}{|x|^2} dx.$$

The problem of positivity of Q_a is solved in the following lemma.

Lemma 2.5. Let $a \in L^\infty(\mathbb{S}^{N-1})$. The following conditions are equivalent:

- i) Q_a is positive definite, i.e. $\inf_{u \in \mathcal{D}^{1,2}(\mathbb{R}^N) \setminus \{0\}} \frac{Q_a(u)}{\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx} > 0$;
- ii) $\Lambda_N(a) < 1$;
- iii) $\mu_1 > -\left(\frac{N-2}{2}\right)^2$ where μ_1 is defined in (9).

Proof. The equivalence between i) and ii) follows from the definition of $\Lambda_N(a)$, see (1). On the other hand, [18, Proposition 1.3 and Lemma 1.1] ensure that i) is equivalent to iii). \square

Remark 1. We notice that in the case of dipole potentials, namely if $a(\theta) = \lambda \frac{x \cdot \mathbf{d}}{|x|}$, then, rewriting (12) in spherical coordinates and exploiting the symmetry with respect to the dipole axis, $\Lambda_N(a)$ can be characterized as

$$\Lambda_N\left(\lambda \frac{x \cdot \mathbf{d}}{|x|}\right) = \lambda \sup_{w \in H_0^1(0, \pi)} \frac{\int_0^\pi \cos \theta w^2(\theta) d\theta}{\int_0^\pi \left[|w'(\theta)|^2 + \frac{(N-2)(N-4)}{4} (\sin \theta)^{-2} w^2(\theta)\right] d\theta}.$$

In dimension $N = 3$, a Taylor's expansion of $\Lambda_N\left(\lambda \frac{x \cdot \mathbf{d}}{|x|}\right)$ near $\lambda = 0$ can be found in [11].

There is no explicit formula for the values $\Lambda_N\left(\lambda \frac{x \cdot \mathbf{d}}{|x|}\right)$. When $N = 4$, it can be expressed in terms of Mathieu's special functions. The results of a numerical approximation of $\Lambda_N\left(\lambda \frac{x \cdot \mathbf{d}}{|x|}\right)$ performed with a finite difference method are listed in table 1, which highlights how the dipole Hardy-type constant detaches the classical Hardy constant more and more as the dimension grows.

N	$\frac{1}{\Lambda_N(1)} = \frac{(N-2)^2}{4}$	$\left[\Lambda_N\left(\frac{x \cdot \mathbf{d}}{ x }\right)\right]^{-1}$	N	$\frac{1}{\Lambda_N(1)} = \frac{(N-2)^2}{4}$	$\left[\Lambda_N\left(\frac{x \cdot \mathbf{d}}{ x }\right)\right]^{-1}$
3	0.25	1.6398	7	6.25	19.0569
4	1	3.7891	8	9	26.7407
5	2.25	7.5831	9	12.25	35.7231
6	4	12.6713	10	16	46.0044

TABLE 1. Some numerical approximations of $\Lambda_N\left(\frac{x \cdot \mathbf{d}}{|x|}\right)$ obtained by finite difference with 10000 steps.

3. A BREZIS-KATO TYPE LEMMA

In this section, we follow the procedure developed by Brezis and Kato in [2] to control from above the behavior of solutions to Schrödinger equations with dipole type potentials, in order to prove Theorem 1.2.

Let us consider the weight φ introduced in (5) and define the weighted H^1 -space $H_\varphi^1(\Omega)$ as the closure of $C^\infty(\bar{\Omega})$ with respect to

$$\|u\|_{H_\varphi^1(\Omega)}^2 := \int_{\Omega} \varphi^2(x) (|\nabla u(x)|^2 + |u(x)|^2) dx,$$

and the space $\mathcal{D}_\varphi^{1,2}(\Omega)$ as the closure of $C_c^\infty(\Omega)$ with respect to

$$\|u\|_{\mathcal{D}_\varphi^{1,2}(\Omega)}^2 := \int_{\Omega} \varphi^2(x) |\nabla u(x)|^2 dx.$$

By the Caffarelli-Kohn-Nirenberg inequality (see [3] and [4]) and the definition of φ , it follows that, for any $w \in \mathcal{D}_\varphi^{1,2}(\Omega)$,

$$(13) \quad \left(\int_{\Omega} \varphi^{2^*}(x) |w(x)|^{2^*} dx \right)^{2/2^*} \leq C_{N,a} \int_{\Omega} \varphi^2(x) |\nabla w|^2 dx,$$

for some positive constant $C_{N,a}$ depending only on N and a .

Lemma 3.1. *Let Ω be a bounded open set containing 0 and $v \in H_\varphi^1(\Omega) \cap L^q(\varphi^{2^*}, \Omega)$, $q > 1$, be a weak solution to*

$$(14) \quad -\operatorname{div}(\varphi^2(x) \nabla v(x)) = \varphi^{2^*}(x) V(x) v(x), \quad \text{in } \Omega,$$

where $V \in L^s(\varphi^{2^*}, \Omega)$ for some $s > N/2$. Then, there exists $\tilde{C} = \tilde{C}(a, N) > 0$ depending only on a and N , such that for any $\Omega' \Subset \Omega$, $v \in L^{\frac{2^*q}{2}}(\varphi^{2^*}, \Omega')$ and

$$(15) \quad \|v\|_{L^{\frac{2^*q}{2}}(\varphi^{2^*}, \Omega')} \leq \tilde{C}^{\frac{1}{q}} (\operatorname{diam} \Omega)^{\frac{\sigma(2-2^*)}{q}} \times \left(\frac{8}{C(q)} \frac{4}{(\operatorname{dist}(\Omega', \partial\Omega))^2} + \frac{4(q+2)}{(\operatorname{dist}(\Omega', \partial\Omega))^2} + \frac{2\ell_q}{C(q)} \right)^{\frac{1}{q}} \|v\|_{L^q(\varphi^{2^*}, \Omega)},$$

where $C(q) := \min \left\{ \frac{1}{4}, \frac{4}{q+4} \right\}$ and

$$\ell_q = \left[\max \left\{ 8C_{N,a} \|V\|_{L^s(\varphi^{2^*}; \Omega)}^{2s/N}, \frac{q+4}{2} C_{N,a} \|V\|_{L^s(\varphi^{2^*}; \Omega)}^{2s/N} \right\} \right]^{\frac{N}{2s-N}}.$$

Proof. Hölder's inequality and (13) yield for any $w \in \mathcal{D}_\varphi^{1,2}(\Omega)$

$$\begin{aligned} & \int_{\Omega} \varphi^{2^*}(x) |V(x)| w^2(x) dx \\ & \leq \ell_q \int_{|V(x)| \leq \ell_q} \varphi^{2^*}(x) w^2(x) dx + \int_{|V(x)| \geq \ell_q} \varphi^{\frac{4}{N-2}}(x) |V(x)| \varphi^2(x) w^2(x) dx \\ (16) \quad & \leq \ell_q \int_{\Omega} \varphi^{2^*}(x) w^2(x) dx + \left[\int_{\Omega} \varphi^{2^*}(x) w^{2^*}(x) dx \right]^{\frac{2}{2^*}} \left[\int_{|V(x)| \geq \ell_q} \varphi^{2^*}(x) |V(x)|^{\frac{N}{2}} dx \right]^{\frac{2}{N}} \\ & \leq \ell_q \int_{\Omega} \varphi^{2^*}(x) w^2(x) dx + C_{N,a} \left[\int_{\Omega} \varphi^2(x) |\nabla w|^2 dx \right] \left[\int_{|V(x)| \geq \ell_q} \varphi^{2^*}(x) |V(x)|^{\frac{N}{2}} dx \right]^{\frac{2}{N}}. \end{aligned}$$

By Hölder's inequality and by the choice of ℓ_q it follows that

$$\begin{aligned} & \int_{|V(x)| \geq \ell_q} \varphi^{2^*}(x) |V(x)|^{\frac{N}{2}} dx \leq \left(\int_{\Omega} \varphi^{2^*}(x) |V(x)|^s dx \right)^{\frac{N}{2s}} \left(\int_{|V(x)| \geq \ell_q} \varphi^{2^*}(x) dx \right)^{\frac{2s-N}{2s}} \\ (17) \quad & \leq \left(\int_{\Omega} \varphi^{2^*}(x) |V(x)|^s dx \right)^{\frac{N}{2s}} \left(\int_{|V(x)| \geq \ell_q} \left(\frac{|V(x)|}{\ell_q} \right)^s \varphi^{2^*}(x) dx \right)^{\frac{2s-N}{2s}} \\ & \leq \|V\|_{L^s(\varphi^{2^*}; \Omega)}^s \ell_q^{-s+\frac{N}{2}} \leq \min \left\{ \frac{C_{N,a}^{-1}}{8}, \frac{2C_{N,a}^{-1}}{q+4} \right\}^{\frac{N}{2}}, \end{aligned}$$

and hence from (16) we obtain that for any $w \in \mathcal{D}_\varphi^{1,2}(\Omega)$

$$\begin{aligned} (18) \quad & \int_{\Omega} \varphi^{2^*}(x) |V(x)| w^2(x) dx \leq \ell_q \int_{\Omega} \varphi^{2^*}(x) w^2(x) dx \\ & \quad + \min \left\{ \frac{1}{8}, \frac{2}{q+4} \right\} \left(\int_{\Omega} \varphi^2(x) |\nabla w(x)|^2 dx \right). \end{aligned}$$

Let η be a nonnegative cut-off function such that

$$\text{supp}(\eta) \Subset \Omega, \quad \eta \equiv 1 \text{ on } \Omega', \quad \text{and } |\nabla \eta(x)| \leq \frac{2}{\text{dist}(\Omega', \partial\Omega)}.$$

Set $v^n := \min(n, |v|) \in \mathcal{D}_\varphi^{1,2}(\Omega)$ and test (14) with $v(v^n)^{q-2}\eta^2 \in \mathcal{D}_\varphi^{1,2}(\Omega)$. This leads to

$$\begin{aligned} (q-2) \int_{\Omega} \varphi^2(x) \eta^2(x) |\nabla v^n(x)|^2 (v^n(x))^{q-2} dx &+ \int_{\Omega} \varphi^2(x) \eta^2(x) (v^n(x))^{q-2} |\nabla v(x)|^2 dx \\ &= \int_{\Omega} \varphi^{2^*}(x) V(x) \eta^2(x) v^2(x) (v^n(x))^{q-2} dx \\ &\quad - 2 \int_{\Omega} \varphi^2(x) \eta(x) v(x) (v^n(x))^{q-2} \nabla v(x) \cdot \nabla \eta(x) dx. \end{aligned}$$

We use the elementary inequality $2ab \leq 1/2a^2 + 4b^2$ and obtain

$$\begin{aligned} (19) \quad &(q-2) \int_{\Omega} \varphi^2(x) \eta^2(x) |\nabla v^n(x)|^2 (v^n(x))^{q-2} dx \\ &\quad + \frac{1}{2} \int_{\Omega} \varphi^2(x) \eta^2(x) (v^n(x))^{q-2} |\nabla v(x)|^2 dx \\ &\leq \int_{\Omega} \varphi^{2^*}(x) V(x) \eta^2(x) v^2(x) (v^n(x))^{q-2} dx + 4 \int_{\Omega} \varphi^2(x) |\nabla \eta(x)|^2 v^2(x) (v^n(x))^{q-2} dx. \end{aligned}$$

Furthermore, an explicit calculation gives

$$\begin{aligned} (20) \quad &|\nabla((v^n)^{\frac{q}{2}-1} v \eta)|^2 \leq \frac{(q+4)(q-2)}{4} (v^n)^{q-2} \eta^2 |\nabla v^n|^2 + 2(v^n)^{q-2} |\nabla v|^2 \eta^2 \\ &\quad + 2(v^n)^{q-2} v^2 |\nabla \eta|^2 + \frac{q-2}{2} (v^n)^q |\nabla \eta|^2. \end{aligned}$$

Letting $C(q) := \min\{\frac{1}{4}, \frac{4}{q+4}\}$, from (19) and (20) we get

$$\begin{aligned} (21) \quad &C(q) \int_{\Omega} \varphi^2(x) |\nabla((v^n)^{\frac{q}{2}-1} v \eta)(x)|^2 dx \\ &\leq 2(2 + C(q)) \int_{\Omega} \varphi^2(x) (v^n(x))^{q-2} v(x)^2 |\nabla \eta(x)|^2 dx \\ &\quad + C(q) \frac{q-2}{2} \int_{\Omega} \varphi^2(x) (v^n(x))^q |\nabla \eta(x)|^2 dx \\ &\quad + \int_{\Omega} \varphi^{2^*}(x) V(x) \eta^2(x) v^2(x) (v^n(x))^{q-2} dx. \end{aligned}$$

Estimate (18) applied to $\eta(v^n)^{\frac{q}{2}-1} v$ gives

$$\begin{aligned} (22) \quad &\int_{\Omega} \varphi^{2^*}(x) |V(x)| [\eta(x) (v^n(x))^{\frac{q}{2}-1} v(x)]^2 dx \\ &\leq \frac{C(q)}{2} \int_{\Omega} \varphi^2(x) |\nabla(\eta(v^n)^{\frac{q}{2}-1} v)(x)|^2 dx + \ell_q \int_{\Omega} \varphi^{2^*}(x) (v^n(x))^{q-2} v^2(x) \eta^2(x) dx. \end{aligned}$$

Using (22) to estimate the term with V in (21), (13) yields

$$\begin{aligned} & \left(\int_{\Omega} \varphi^{2^*}(x) |v^n(x)|^{(\frac{q}{2}-1)2^*} |v(x)|^{2^*} \eta^{2^*}(x) dx \right)^{\frac{2}{2^*}} \\ & \leq \frac{2\ell_q C_{N,a}}{C(q)} \int_{\Omega} \varphi^{2^*}(x) \eta^2(x) |v^n(x)|^{q-2} v^2(x) dx \\ & \quad + \frac{4C_{N,a}(2+C(q))}{C(q)} \int_{\Omega} \varphi^2(x) |v^n(x)|^{q-2} v^2(x) |\nabla \eta(x)|^2 dx \\ & \quad + C_{N,a}(q-2) \int_{\Omega} \varphi^2(x) |v^n(x)|^q |\nabla \eta(x)|^2 dx. \end{aligned}$$

Letting $n \rightarrow \infty$ in the above inequality, (15) follows. \square

Proof of Theorem 1.2. Let u be a weak $H^1(\Omega)$ -solution to (6). It is easy to verify that $\varphi(x) := |x|^\sigma \psi_1(x/|x|) \in H^1(\Omega)$ satisfies (in a weak $H^1(\Omega)$ -sense and in a classical sense in $\Omega \setminus \{0\}$)

$$-\Delta \varphi(x) - \frac{a(x/|x|)}{|x|^2} \varphi(x) = 0.$$

Then $v := \frac{u}{\varphi} \in H_\varphi^1(\Omega)$ turns out to be a weak solution to (14). Let $R > 0$ be such that

$$\Omega' \Subset \Omega' + B(0, 2R) \Subset \Omega.$$

Using Lemma 3.1 in $\Omega_1 := \Omega' + B(0, R(2-r_1)) \Subset \Omega' + B(0, 2R)$, $r_1 = 1$, with $q = q_1 = 2^*$, we infer that $v \in L^{\frac{(2^*)^2}{2}}(\varphi^{2^*}, \Omega_1)$ and the following estimate holds

$$\begin{aligned} & \|v\|_{L^{\frac{(2^*)^2}{2}}(\varphi^{2^*}, \Omega_1)} \\ & \leq \tilde{C}^{\frac{1}{q_1}} (\text{diam } \Omega)^{\frac{\sigma(2-2^*)}{q_1}} \left(\frac{8}{C(q_1)} \frac{4}{(Rr_1)^2} + \frac{4(q_1+2)}{(Rr_1)^2} + \frac{2\ell_{q_1}}{C(q_1)} \right)^{\frac{1}{q_1}} \|v\|_{L^{2^*}(\varphi^{2^*}, \Omega)}. \end{aligned}$$

Using again Lemma 3.1 in $\Omega_2 := \Omega' + B(0, R(2-r_1-r_2)) \Subset \Omega_1$, $r_2 = \frac{1}{4}$, with $q = q_2 = (2^*)^2/2$, we infer that $v \in L^{\frac{(2^*)^3}{4}}(\varphi^{2^*}, \Omega_2)$ and

$$\begin{aligned} & \|v\|_{L^{\frac{(2^*)^3}{4}}(\varphi^{2^*}, \Omega_2)} \\ & \leq \tilde{C}^{\frac{1}{q_2}} (\text{diam } \Omega)^{\frac{\sigma(2-2^*)}{q_2}} \left(\frac{8}{C(q_2)} \frac{4}{(Rr_2)^2} + \frac{4(q_2+2)}{(Rr_2)^2} + \frac{2\ell_{q_2}}{C(q_2)} \right)^{\frac{1}{q_2}} \|v\|_{L^{q_2}(\varphi^{2^*}, \Omega_1)} \\ & \leq \left[\tilde{C} (\text{diam } \Omega)^{\sigma(2-2^*)} \right]^{\frac{1}{q_1} + \frac{1}{q_2}} \left(\frac{8}{C(q_1)} \frac{4}{(Rr_1)^2} + \frac{4(q_1+2)}{(Rr_1)^2} + \frac{2\ell_{q_1}}{C(q_1)} \right)^{\frac{1}{q_1}} \times \\ & \quad \times \left(\frac{8}{C(q_2)} \frac{4}{(Rr_2)^2} + \frac{4(q_2+2)}{(Rr_2)^2} + \frac{2\ell_{q_2}}{C(q_2)} \right)^{\frac{1}{q_2}} \|v\|_{L^{2^*}(\varphi^{2^*}, \Omega)}. \end{aligned}$$

Setting, for any $n \in \mathbb{N}$, $n \geq 1$,

$$q_n = \frac{1}{2} \left(\frac{2^*}{2} \right)^n, \quad \Omega_n := \Omega' + B\left(0, R\left(2 - \sum_{k=1}^n r_k\right)\right), \quad \text{and} \quad r_n = \frac{1}{n^2},$$

and using iteratively Lemma 3.1, we obtain that, for any $n \in \mathbb{N}$, $n \geq 1$,

$$(23) \quad \begin{aligned} \|v\|_{L^{q_{n+1}}(\varphi^{2^*}, \Omega')} &\leq \|v\|_{L^{q_{n+1}}(\varphi^{2^*}, \Omega_n)} \\ &\leq \|v\|_{L^{2^*}(\varphi^{2^*}, \Omega)} \left[\tilde{C} (\text{diam } \Omega)^{\sigma(2-2^*)} \right]^{\sum_{k=1}^n \frac{1}{q_k}} \times \\ &\quad \times \prod_{k=1}^n \left(\frac{8}{C(q_k)} \frac{4}{(Rr_k)^2} + \frac{4(q_k+2)}{(Rr_k)^2} + \frac{2\ell_{q_k}}{C(q_k)} \right)^{\frac{1}{q_k}}. \end{aligned}$$

We notice that

$$(24) \quad \prod_{k=1}^n \left(\frac{8}{C(q_k)} \frac{4}{(Rr_k)^2} + \frac{4(q_k+2)}{(Rr_k)^2} + \frac{2\ell_{q_k}}{C(q_k)} \right)^{\frac{1}{q_k}} = \exp \left[\sum_{k=1}^n b_k \right]$$

where

$$b_k = \frac{1}{q_k} \log \left(\frac{16k^4}{R^2 C(q_k)} + \frac{4k^4(q_k+2)}{R^2} + \frac{2\ell_{q_k}}{C(q_k)} \right),$$

and, for some constant $C = C(N, a, \|V\|_{L^s(\varphi^{2^*}, \Omega)}, \text{dist}(\Omega', \partial\Omega)) > 0$,

$$b_k \sim 2 \left(\frac{2}{2^*} \right)^k \log \left[C \left(\frac{1}{2} \left(\frac{2^*}{2} \right) \right)^{\frac{2s}{2s-N}} \right] \quad \text{as } k \rightarrow +\infty.$$

Hence $\sum_{n=1}^{\infty} b_n$ converges to some positive sum depending only on $\|V\|_{L^s(\varphi^{2^*}, \Omega)}$, $\text{dist}(\Omega', \partial\Omega)$, N , and a , hence

$$\lim_{n \rightarrow +\infty} \left[\tilde{C} (\text{diam } \Omega)^{\sigma(2-2^*)} \right]^{\sum_{k=1}^n \frac{1}{q_k}} \prod_{k=1}^n \left(\frac{8}{C(q_k)} \frac{4}{(Rr_k)^2} + \frac{4(q_k+2)}{(Rr_k)^2} + \frac{2\ell_{q_k}}{C(q_k)} \right)^{\frac{1}{q_k}}$$

is finite and depends only on N , a , $\|V\|_{L^s(\varphi^{2^*}, \Omega)}$, and $\text{dist}(\Omega', \partial\Omega)$. Hence, from (23), we deduce that there exists a positive constant C depending only on N , a , $\|V\|_{L^s(\varphi^{2^*}, \Omega)}$, $\text{dist}(\Omega', \partial\Omega)$, and $\text{diam } \Omega$, such that

$$\|v\|_{L^{q_{n+1}}(\varphi^{2^*}, \Omega')} \leq C \|v\|_{L^{2^*}(\varphi^{2^*}, \Omega)} \quad \text{for all } n \in \mathbb{N}.$$

Letting $n \rightarrow +\infty$ we deduce that v is essentially bounded in Ω' with respect to the measure $\varphi^{2^*} dx$ and

$$\|v\|_{L^\infty(\varphi^{2^*}, \Omega')} \leq C \|v\|_{L^{2^*}(\varphi^{2^*}, \Omega)} = C \|u\|_{L^{2^*}(\Omega)},$$

where $\|v\|_{L^\infty(\varphi^{2^*}, \Omega')}$ denotes the essential supremum of v with respect to the measure $\varphi^{2^*} dx$. Since $\varphi^{2^*} dx$ is absolutely continuous with respect to the Lebesgue measure and viceversa, there holds $\|v\|_{L^\infty(\varphi^{2^*}, \Omega')} = \|v\|_{L^\infty(\Omega')}$, hence $v \in L^\infty(\Omega')$ and

$$\|v\|_{L^\infty(\Omega')} \leq C \|u\|_{L^{2^*}(\Omega)},$$

thus completing the proof. \square

If the potential V in equation (6) belongs to $L^{N/2}(\varphi^{2^*}, \Omega)$ (but to $L^s(\varphi^{2^*}, \Omega)$ for no $s > N/2$), although we can no more derive an L^∞ -bound for u/φ , we can obtain for u/φ as high summability as we like.

Theorem 3.2. *Let Ω be a bounded domain containing 0, $a \in L^\infty(\mathbb{S}^{N-1})$ satisfying $\Lambda_N(a) < 1$, and $V \in L^{N/2}(\varphi^{2^*}, \Omega)$. Then, for any $\Omega' \Subset \Omega$ and for any weak $H^1(\Omega)$ -solution u to (6), there holds $\frac{u}{\varphi} \in L^q(\varphi^{2^*}, \Omega')$ for all $1 \leq q < +\infty$.*

Proof. The proof follows closely the proofs of Theorem 1.2 and Lemma 3.1. However, since we only require $V \in L^{N/2}(\varphi^{2^*}, \Omega)$, we have that for any q there exists ℓ_q such that

$$\int_{|V(x)| \geq \ell_q} \varphi^{2^*}(x) |V(x)|^{\frac{N}{2}} dx \leq \min \left\{ \frac{C_{N,a}^{-1}}{8}, \frac{2C_{N,a}^{-1}}{q+4} \right\}^{\frac{N}{2}},$$

but we can no more estimate ℓ_q in terms of q , as we did in (17) thanks to the summability assumption $V \in L^s(\varphi^{2^*}, \Omega)$ for some $s > N/2$. Hence we still arrive at an estimate of type (23) but we have no control on the product in (24) as $n \rightarrow +\infty$. \square

4. BEHAVIOR OF SOLUTIONS AT SINGULARITIES

The procedure followed in this section to prove Theorem 1.1 relies in comparison methods and separation of variables. Indeed we will evaluate the asymptotics of solutions to problem (3) by trapping them between functions which solve analogous problems with radial perturbing potentials. To this aim, the first step consists in deriving the asymptotic behavior of solutions to Schrödinger equations with a potential which is given by a radial perturbation of the dipole-type singular term. In this case, it is possible to expand the solution in Fourier series, thus separating the radial and angular variables, and to estimate the behavior of the Fourier coefficients in order to establish which of them is dominant near the singularity.

Proposition 1. *Let $a \in L^\infty(\mathbb{S}^{N-1})$ be such that $\Lambda_N(a) < 1$ and $R > 0$. Assume that $u \in H^1(B(0, R))$, $u \geq 0$ a.e. in $B(0, R)$, $u \not\equiv 0$, is a weak H^1 -solution to*

$$(25) \quad -\Delta u(x) = \left[\frac{a(x/|x|)}{|x|^2} + h(|x|) \right] u(x) \quad \text{in } B(0, R),$$

where $h \in L^\infty_{\text{loc}}(0, R) \cap L^p(0, R)$ for some $p > N/2$. Then, for any $r \in (0, R)$, there exists a positive constant C (depending on h, R, r, a, ε , and u) such that

$$\frac{1}{C} |x|^\sigma \leq u(x) \leq C |x|^\sigma \quad \text{for all } x \in B(0, r) \setminus \{0\},$$

where σ is defined in (2). Moreover, there exists a positive constant \tilde{C} (depending on u, h, N , and a) such that, for any $\theta \in \mathbb{S}^{N-1}$,

$$(26) \quad \lim_{\rho \rightarrow 0^+} u(\rho \theta) \rho^{-\sigma} = \tilde{C} \psi_1(\theta),$$

and, for any $r \in (0, R)$,

$$(27) \quad \tilde{C} = \int_{\mathbb{S}^{N-1}} \left(r^{-\sigma} u(r\eta) + \int_0^r \frac{s^{1-\sigma}}{2\sigma + N - 2} h(s) u(s\eta) ds \right. \\ \left. - r^{-2\sigma - N + 2} \int_0^r \frac{s^{N-1+\sigma}}{2\sigma + N - 2} h(s) u(s\eta) ds \right) \psi_1(\eta) dV(\eta).$$

Furthermore, for any $r \in (0, R)$, there exists a positive constant \bar{C} (depending on h, R, r, a, ε , but not on u) such that

$$(28) \quad u(\rho\theta) \leq \bar{C} \|u\|_{H^1(B(0,R))} \rho^\sigma, \quad \text{for all } 0 < \rho < r.$$

Proof. Let $r \in (0, R)$. We can assume, without loss of generality, that $R > 1$ and $r = 1$. Indeed, setting $w(x) := u(rx)$, we notice that $w \in H^1(B(0, R/r))$ and weakly solves

$$-\Delta w(x) = \left[\frac{a(x/|x|)}{|x|^2} + \tilde{h}(|x|) \right] w(x) \quad \text{in } B(0, R/r),$$

where $\tilde{h}(\rho) := r^2 h(r\rho)$ satisfies $\tilde{h} \in L_{\text{loc}}^\infty(0, R/r) \cap L^p(0, R/r)$. Hence, it is enough to prove the statement for $R > 1$ and $r = 1$, being the general case easily obtainable from scaling.

Let $R > 1, r = 1$ and $u \in H^1(B(0, R))$, $u \geq 0$ a.e. in $B(0, R)$, $u \not\equiv 0$, be a weak solution of (25). By standard regularity theory, $u \in C^0(\bar{B}(0, 1) \setminus B(0, s))$ for any $s \in (0, 1)$. For any $k \in \mathbb{N} \setminus \{0\}$, let ψ_k be a L^2 -normalized eigenfunction of the operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ on the sphere associated to the k -th eigenvalue μ_k , i.e. satisfying (11). We can choose the functions ψ_k in such a way that they form an orthonormal basis of $L^2(\mathbb{S}^{N-1})$, hence u can be expanded as

$$(29) \quad u(x) = u(\rho\theta) = \sum_{k=1}^{\infty} \varphi_k(\rho) \psi_k(\theta),$$

where $\rho = |x| \in (0, 1]$, $\theta = x/|x| \in \mathbb{S}^{N-1}$, and

$$(30) \quad \varphi_k(\rho) = \int_{\mathbb{S}^{N-1}} u(\rho\theta) \psi_k(\theta) dV(\theta).$$

The Parseval identity yields

$$\int_{\mathbb{S}^{N-1}} |u(\rho\theta)|^2 dV(\theta) = \sum_{k=1}^{\infty} |\varphi_k(\rho)|^2, \quad \text{for all } 0 < \rho \leq 1,$$

and hence

$$(31) \quad \|u\|_{L^2(B(0,1))}^2 = \int_0^1 \rho^{N-1} \left(\sum_{k=1}^{\infty} |\varphi_k(\rho)|^2 \right) d\rho = \sum_{k=1}^{\infty} \int_0^1 \rho^{N-1} |\varphi_k(\rho)|^2 d\rho.$$

Equations (25) and (11) imply that, for every k ,

$$\varphi_k''(\rho) + \frac{N-1}{\rho} \varphi_k'(\rho) - \frac{\mu_k}{\rho^2} \varphi_k(\rho) = h(\rho) \varphi_k(\rho), \quad \text{in } (0, 1).$$

A direct calculation shows that, for some $c_1^k, c_2^k \in \mathbb{R}$,

$$(32) \quad \begin{aligned} \varphi_k(\rho) = & \rho^{\sigma_k^+} \left(c_1^k + \int_{\rho}^1 \frac{s^{-\sigma_k^+ + 1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds \right) \\ & + \rho^{\sigma_k^-} \left(c_2^k + \int_{\rho}^1 \frac{s^{-\sigma_k^- + 1}}{\sigma_k^- - \sigma_k^+} h(s) \varphi_k(s) ds \right), \end{aligned}$$

where

$$(33) \quad \sigma_k^+ = -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k} \quad \text{and} \quad \sigma_k^- = -\frac{N-2}{2} - \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k}.$$

For the sake of notation, we set

$$(34) \quad A_k(\rho) = \rho^{\sigma_k^+} \int_{\rho}^1 \frac{s^{-\sigma_k^++1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds$$

and

$$B_k(\rho) = \rho^{\sigma_k^-} \left(c_2^k + \int_{\rho}^1 \frac{s^{-\sigma_k^-+1}}{\sigma_k^- - \sigma_k^+} h(s) \varphi_k(s) ds \right),$$

so that

$$(35) \quad \varphi_k(\rho) = c_1^k \rho^{\sigma_k^+} + A_k(\rho) + B_k(\rho).$$

Without loss of generality, we can assume that

$$\frac{N}{2} < p < \frac{N}{2 - \frac{1}{3} \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1}},$$

so, setting $\varepsilon = 2 - \frac{N}{p}$, $0 < \varepsilon < \frac{1}{3}(\sigma_k^+ + \frac{N-2}{2})$, for every k . From Hölder's inequality and Lemma 2.3, it follows that

$$(36) \quad \begin{aligned} |A_k(\rho)| &= \rho^{\sigma_k^+} \left| \int_{\rho}^1 \frac{s^{-\sigma_k^++1}}{\sigma_k^+ - \sigma_k^-} h(s) \left(\int_{\mathbb{S}^{N-1}} u(s\theta) \psi_k(\theta) dV(\theta) \right) ds \right| \\ &\leq \frac{C_1 \rho^{\sigma_k^+} |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1}}{\sigma_k^+ - \sigma_k^-} \times \\ &\quad \times \int_{\rho}^1 s^{-\sigma_k^++1 - \frac{N-1}{p} - \frac{N-1}{2^*}} \left(\int_{\mathbb{S}^{N-1}} s^{\frac{N-1}{p} + \frac{N-1}{2^*}} |h(s)| |u(s\theta)| dV(\theta) \right) ds \\ &\leq \frac{\omega_N^{\frac{N+2}{2N}} C_1 \rho^{\sigma_k^+} |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1}}{\sigma_k^+ - \sigma_k^-} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \times \\ &\quad \times \left(\int_{\rho}^1 s^{(-\sigma_k^++1 - \frac{N-1}{p} - \frac{N-1}{2^*}) \frac{2^* p}{2^* p - 2^* - p}} ds \right)^{1 - \frac{1}{p} - \frac{1}{2^*}} \\ &= \frac{\omega_N^{\frac{N+2}{2N}} C_1 \rho^{\sigma_k^+} |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1}}{\sigma_k^+ - \sigma_k^-} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \times \\ &\quad \times \left[\frac{\rho^{\frac{2^* p}{2^* p - 2^* - p} (-\sigma_k^+ - \frac{N-2}{2} + 2 - \frac{N}{p})}}{\frac{2^* p}{2^* p - 2^* - p} \left(\frac{N}{p} - 2 + \sigma_k^+ + \frac{N-2}{2} \right)} \right]^{1 - \frac{1}{p} - \frac{1}{2^*}} \\ &\leq \frac{\omega_N^{\frac{N+2}{2N}} C_1 |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1}}{\sigma_k^+ - \sigma_k^-} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \times \\ &\quad \times \frac{\rho^{\varepsilon - \frac{N-2}{2}}}{\left[\frac{2^* p}{2^* p - 2^* - p} \left(\sigma_k^+ + \frac{N-2}{2} - \varepsilon \right) \right]^{1 - \frac{1}{p} - \frac{1}{2^*}}}, \end{aligned}$$

where $\omega_N = \int_{\mathbb{S}^{N-1}} dV(\theta)$. In particular

$$(37) \quad A_k(\rho) = o(\rho^{\sigma_k^-}) \quad \text{as } \rho \rightarrow 0^+.$$

Moreover

$$\begin{aligned}
& \int_0^1 |s^{-\sigma_k^-+1} h(s) \varphi_k(s)| ds \\
& \leq C_1 |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \times \\
& \quad \times \int_0^1 s^{-\sigma_k^-+1 - \frac{N-1}{p} - \frac{N-1}{2^*}} \left(\int_{\mathbb{S}^{N-1}} s^{\frac{N-1}{p} + \frac{N-1}{2^*}} |h(s)| |u(s\theta)| dV(\theta) \right) ds \\
(38) \quad & \leq \omega_N^{\frac{N+2}{2N}} C_1 |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \times \\
& \quad \times \left[\int_0^1 s^{(-\sigma_k^-+1 - \frac{N-1}{p} - \frac{N-1}{2^*}) \frac{2^*p}{2^*p-2^*-p}} ds \right]^{1-\frac{1}{p}-\frac{1}{2^*}} \\
& = \omega_N^{\frac{N+2}{2N}} C_1 |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \times \\
& \quad \times \frac{1}{\left[\frac{2^*p}{2^*p-2^*-p} \left(\varepsilon - \sigma_k^- - \frac{N-2}{2} \right) \right]^{1-\frac{1}{p}-\frac{1}{2^*}}} < \infty
\end{aligned}$$

due to inequality $\sigma_k^- < -\frac{N-2}{2}$. As a consequence,

$$(39) \quad \lim_{\rho \rightarrow 0^+} \int_\rho^1 \frac{s^{-\sigma_k^-+1}}{\sigma_k^- - \sigma_k^+} h(s) \varphi_k(s) ds$$

is finite. Since $u \in L^{2^*}(B(0,1))$, from (37), (39), and the fact that $\rho^{\sigma_k^-} \psi_k(\theta) \notin L^{2^*}(B(0,1))$, we conclude that there must be

$$(40) \quad c_2^k = - \int_0^1 \frac{s^{-\sigma_k^-+1}}{\sigma_k^- - \sigma_k^+} h(s) \varphi_k(s) ds,$$

hence

$$(41) \quad B_k(\rho) = \rho^{\sigma_k^-} \int_0^\rho \frac{s^{-\sigma_k^-+1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds.$$

Since $u \in C^0(\overline{B(0,1)} \setminus B(0,s))$ for any $s \in (0,1)$, it makes sense to evaluate φ_k at $\rho = 1$ and, from (32) and (40), we have that

$$(42) \quad c_1^k = \varphi_k(1) + \int_0^1 \frac{s^{-\sigma_k^-+1}}{\sigma_k^- - \sigma_k^+} h(s) \varphi_k(s) ds.$$

From (32), (41), and (42), we deduce that

$$\begin{aligned}
(43) \quad \varphi_k(\rho) & = \rho^{\sigma_k^+} \left(\varphi_k(1) + \int_0^1 \frac{s^{-\sigma_k^-+1}}{\sigma_k^- - \sigma_k^+} h(s) \varphi_k(s) ds \right. \\
& \quad \left. + \int_\rho^1 \frac{s^{-\sigma_k^++1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds \right) + \rho^{\sigma_k^-} \int_0^\rho \frac{s^{-\sigma_k^-+1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds.
\end{aligned}$$

From above, (38), and standard elliptic estimates (which allow to estimate u outside the singularity in terms of its H^1 -norm) we obtain that, for some positive constant \tilde{c} depending only on $N, R, a,$

and h ,

$$(44) \quad |c_1^k| \leq \tilde{c} |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{H^1(B(0,R))} \left[1 + \left[\frac{2^* p}{2^* p - 2^* - p} (\varepsilon - \sigma_k^- - \frac{N-2}{2}) \right]^{-1 + \frac{1}{p} + \frac{1}{2^*}} \right].$$

Arguing as in (38), we find that

$$(45) \quad |B_k(\rho)| \leq \frac{\omega_N^{\frac{N+2}{2N}} C_1 |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|h\|_{L^p(B(0,1))}}{\sigma_k^+ - \sigma_k^-} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \times \frac{\rho^{\varepsilon - \frac{N-2}{2}}}{\left[\frac{2^* p}{2^* p - 2^* - p} (\varepsilon - \sigma_k^- - \frac{N-2}{2}) \right]^{1 - \frac{1}{p} - \frac{1}{2^*}}}.$$

From (35), (36), and (45), we can estimate φ_k as

$$(46) \quad |\varphi_k(\rho)| \leq |c_1^k| \rho^{\sigma_k^+} + \frac{\alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|h\|_{L^p(B(0,1))}}{\sigma_k^+ - \sigma_k^-} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N} \|u\|_{L^{2^*}(B(0,1))} \rho^{\varepsilon - \frac{N-2}{2}}$$

where

$$\alpha_k := \frac{\omega_N^{\frac{N+2}{2N}} C_1}{\left[\frac{2^* p}{2^* p - 2^* - p} (\sigma_k^+ + \frac{N-2}{2} - \varepsilon) \right]^{1 - \frac{1}{p} - \frac{1}{2^*}}} + \frac{\omega_N^{\frac{N+2}{2N}} C_1}{\left[\frac{2^* p}{2^* p - 2^* - p} (\varepsilon - \sigma_k^- - \frac{N-2}{2}) \right]^{1 - \frac{1}{p} - \frac{1}{2^*}}}.$$

Claim 1: there holds

$$(47) \quad |A_k(\rho) + B_k(\rho)| \leq |c_1^k| \rho^{\sigma_k^+} \sum_{i=1}^{j_k-1} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-)} \frac{2}{\varepsilon^{\frac{p-1}{p}}} \right)^i + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-)} \right)^{j_k} \times \rho^{j_k \varepsilon - \frac{N-2}{2}} \prod_{i=2}^{j_k} \frac{2}{\left[\sqrt{\left(\frac{N-2}{2} \right)^2 + \mu_k - i\varepsilon} \right]^{\frac{p-1}{p}}},$$

where

$$j_k := \left\lfloor \frac{1}{\varepsilon} \sqrt{\left(\frac{N-2}{2} \right)^2 + \mu_k} \right\rfloor - 1,$$

i.e. j_k is the unique integer number such that

$$\sqrt{\left(\frac{N-2}{2} \right)^2 + \mu_k - 2\varepsilon} < j_k \varepsilon \leq \sqrt{\left(\frac{N-2}{2} \right)^2 + \mu_k} - \varepsilon.$$

Notice that $\varepsilon \leq \frac{1}{3} (\sigma_k^+ + \frac{N-2}{2})$ implies $j_k \geq 2$.

To prove the claim, we observe that, from (34) and (46) it follows that

$$\begin{aligned}
|A_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \left(\int_\rho^1 s^{(1-\frac{N-1}{p})\frac{p}{p-1}} ds \right)^{\frac{p-1}{p}} \\
&\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^2 \times \\
&\quad \quad \quad \times \rho^{\sigma_k^+} \left(\int_\rho^1 s^{(-\sigma_k^+ + 1 + \varepsilon - \frac{N-2}{2} - \frac{N-1}{p})\frac{p}{p-1}} ds \right)^{\frac{p-1}{p}} \\
(48) \quad &\leq |c_1^k| \rho^{\sigma_k^+} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{1}{\left(\varepsilon \frac{p}{p-1}\right)^{\frac{p-1}{p}}} + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \times \\
&\quad \quad \quad \times \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^2 \frac{\rho^{2\varepsilon - \frac{N-2}{2}}}{\left[(\sigma_k^+ - 2\varepsilon + \frac{N-2}{2}) \frac{p}{p-1} \right]^{\frac{p-1}{p}}}.
\end{aligned}$$

In a similar way, from (41) and (46) we deduce that

$$\begin{aligned}
|B_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{1}{\left[(\varepsilon + \sigma_k^+ - \sigma_k^-) \frac{p}{p-1} \right]^{\frac{p-1}{p}}} \\
(49) \quad &\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left[\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right]^2 \frac{\rho^{2\varepsilon - \frac{N-2}{2}}}{\left[(2\varepsilon - \sigma_k^- - \frac{N-2}{2}) \frac{p}{p-1} \right]^{\frac{p-1}{p}}}.
\end{aligned}$$

Summing up (48) and (49), we obtain

$$\begin{aligned}
|A_k(\rho) + B_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{2}{\varepsilon \frac{p}{p-1}} \\
&\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left[\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right]^2 \frac{2\rho^{2\varepsilon - \frac{N-2}{2}}}{\left(\sigma_k^+ + \frac{N-2}{2} - 2\varepsilon \right)^{\frac{p-1}{p}}},
\end{aligned}$$

and hence, from (35),

$$\begin{aligned}
|\varphi_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \left(1 + \frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{2}{\varepsilon \frac{p}{p-1}} \right) \\
(50) \quad &\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left[\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right]^2 \frac{2\rho^{2\varepsilon - \frac{N-2}{2}}}{\left(\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k} - 2\varepsilon \right)^{\frac{p-1}{p}}}.
\end{aligned}$$

Using (50), we can improve our estimates of $A_k(\rho)$ and $B_k(\rho)$ thus obtaining

$$\begin{aligned} |A_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \left[\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{1}{\varepsilon^{\frac{p-1}{p}}} + \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^2 \frac{2}{\varepsilon^{2\frac{p-1}{p}}} \right] \\ &\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^3 \times \\ &\quad \times \frac{\rho^{3\varepsilon - \frac{N-2}{2}}}{\left[\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k - 3\varepsilon} \right]^{\frac{p-1}{p}}} \frac{2}{\left[\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k - 2\varepsilon} \right]^{\frac{p-1}{p}}} \end{aligned}$$

and

$$\begin{aligned} |B_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \left[\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{1}{\varepsilon^{\frac{p-1}{p}}} + \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^2 \frac{2}{\varepsilon^{2\frac{p-1}{p}}} \right] \\ &\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^3 \times \\ &\quad \times \frac{\rho^{3\varepsilon - \frac{N-2}{2}}}{\left[\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k + 3\varepsilon} \right]^{\frac{p-1}{p}}} \frac{2}{\left[\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k - 2\varepsilon} \right]^{\frac{p-1}{p}}}. \end{aligned}$$

Summing up we find that

$$\begin{aligned} |A_k(\rho) + B_k(\rho)| &\leq |c_1^k| \rho^{\sigma_k^+} \left[\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \frac{2}{\varepsilon^{\frac{p-1}{p}}} + \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^2 \frac{4}{\varepsilon^{2\frac{p-1}{p}}} \right] \\ &\quad + \alpha_k |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \|u\|_{L^{2^*}(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N(\sigma_k^+ - \sigma_k^-)} \right)^3 \times \\ &\quad \times \frac{4\rho^{3\varepsilon - \frac{N-2}{2}}}{\left[\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k - 3\varepsilon} \right]^{\frac{p-1}{p}} \left[\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k - 2\varepsilon} \right]^{\frac{p-1}{p}}}. \end{aligned}$$

An iteration of the above argument ($j_k - 1$) times easily leads to estimate (47). Claim 1 is thereby proved.

Claim 2: the function $s \mapsto s^{-\sigma_k^+ + 1} h(s) \varphi_k(s)$ belongs to $L^1(0, 1)$ and

$$(51) \quad \lim_{\rho \rightarrow 0^+} \rho^{-\sigma_k^+} \varphi_k(\rho) = c_1^k + \int_0^1 \frac{s^{-\sigma_k^+ + 1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds.$$

Indeed, from (47), (35), (44), and the choice of j_k , it follows that

$$(52) \quad |\varphi_k(\rho)| \leq d_k \|u\|_{H^1(B(0,1))} \rho^{j_k \varepsilon - \frac{N-2}{2}},$$

for some positive constant d_k depending on k (and on a, R, h, N). We distinguish now two cases.

If $j_k \varepsilon < \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k} - \varepsilon$, then from (34), (41), and (52) we derive that

$$|A_k(\rho) + B_k(\rho)| \leq d'_k \|u\|_{H^1(B(0,1))} \rho^{(j_k + 1)\varepsilon - \frac{N-2}{2}},$$

for some other positive constant d'_k depending on k (and on a, R, h, N), and hence, by (44) and the choice of j_k ,

$$(53) \quad |\varphi_k(\rho)| \leq d'_k \|u\|_{H^1(B(0,1))} \rho^{(j_k+1)\varepsilon - \frac{N-2}{2}}.$$

Estimate (53) and the choice of j_k imply that the function $s \mapsto s^{-\sigma_k^+ + 1} h(s) \varphi_k(s)$ belongs to $L^1(0,1)$. Moreover, from (41) and (53) it follows that

$$(54) \quad |B_k(\rho)| \leq d''_k \|u\|_{H^1(B(0,1))} \rho^{(j_k+2)\varepsilon - \frac{N-2}{2}} = o(\rho^{\sigma_k^+}) \quad \text{as } \rho \rightarrow 0^+,$$

and the claim is proved. If $j_k \varepsilon = \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k} - \varepsilon$, then from (34), (41), and (52) we derive that

$$|A_k(\rho) + B_k(\rho)| \leq \gamma_k \|u\|_{H^1(B(0,1))} \rho^{\sigma_k^+} |\log \rho|^{\frac{p-1}{p}},$$

for some other positive constant γ_k depending on k (and on a, R, h, N), and hence,

$$(55) \quad |\varphi_k(\rho)| \leq \gamma'_k \|u\|_{H^1(B(0,1))} \rho^{\sigma_k^+} |\log \rho|^{\frac{p-1}{p}}.$$

Estimate (55) implies that the function $s \mapsto s^{-\sigma_k^+ + 1} h(s) \varphi_k(s)$ belongs to $L^1(0,1)$. Moreover, from (41) and (55) it follows that

$$(56) \quad |B_k(\rho)| \leq \gamma''_k \|u\|_{H^1(B(0,1))} \rho^{\sigma_k^-} \left(\int_0^\rho s^{(\sigma_k^+ - \sigma_k^- + \varepsilon) \frac{p}{p-1} - 1} |\log s| ds \right)^{\frac{p-1}{p}} = o(\rho^{\sigma_k^+})$$

as $\rho \rightarrow 0^+$, and claim 2 is proved also in this case.

Let us fix \bar{k} such that

$$(57) \quad \frac{\|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-)} \frac{2}{\varepsilon^{\frac{p-1}{p}}} = \frac{\|h\|_{L^p(B(0,1))}}{\omega_N \varepsilon^{\frac{p-1}{p}} \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k}} < \frac{1}{3},$$

$$\sigma_k^+ > 4\varepsilon \quad \text{and} \quad \frac{\sigma_k^+}{2} > \sigma_1^+, \quad \forall k \geq \bar{k}.$$

From (47) and (57), it follows that, for all $k \geq \bar{k}$ and for some positive constant C_2 (depending only on N, h , and a),

$$|A_k(\rho) + B_k(\rho)| \leq |c_1^k| \rho^{\sigma_k^+} \sum_{i=1}^{\infty} \left(\frac{1}{3}\right)^i + C_2 \|u\|_{H^1(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-)} \right)^{j_k} \times$$

$$\times |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \frac{\rho^{j_k \varepsilon - \frac{N-2}{2}} \left(\frac{2}{\varepsilon}\right)^{j_k - 1}}{\prod_{i=2}^{j_k} \left(\left[\varepsilon^{-1} \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k} \right] - i \right)},$$

which yields, for all $k \geq \bar{k}$,

$$(58) \quad |A_k(\rho) + B_k(\rho)| \leq \frac{1}{2} |c_1^k| \rho^{\sigma_k^+} + b_k \rho^{\sigma_k^+ / 2},$$

where

$$b_k = C_2 \|u\|_{H^1(B(0,1))} \left(\frac{\|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-)} \right)^{j_k} |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \frac{\left(\frac{2}{\varepsilon}\right)^{j_k - 1}}{(j_k - 1)!}.$$

From (10), $\mu_k \sim k^{2/(N-1)}$ and $j_k \sim k^{1/(N-1)}$ as $k \rightarrow +\infty$. Hence we have that

$$(59) \quad |b_k| \leq C_3 \|u\|_{H^1(B(0,1))} \exp(-C_4 k^{1/(N-1)}),$$

for some positive constants C_3 and C_4 depending only on N , h , and a . In view of (35) and (58), we deduce that

$$\frac{1}{2} |c_1^k| \rho^{\sigma_k^+} \leq |\varphi_k(\rho)| + |b_k| \rho^{\sigma_k^+/2}, \quad \text{for all } k \geq \bar{k},$$

and consequently

$$\begin{aligned} & \frac{1}{4} \sum_{k=\bar{k}}^{\infty} |c_1^k|^2 \int_0^1 \rho^{2\sigma_k^+ + N-1} d\rho \\ & \leq 2 \sum_{k=\bar{k}}^{\infty} \int_0^1 |\varphi_k(\rho)|^2 \rho^{N-1} d\rho + 2 \sum_{k=\bar{k}}^{\infty} |b_k|^2 \int_0^1 \rho^{\sigma_k^+ + N-1} d\rho. \end{aligned}$$

Hence, from (31) and (59), we obtain that

$$(60) \quad \sum_{k=\bar{k}}^{\infty} \frac{|c_1^k|^2}{N + 2\sigma_k^+} \leq 8 \|u\|_{L^2(B(0,1))}^2 + 8 \sum_{k=\bar{k}}^{\infty} \frac{|b_k|^2}{N + \sigma_k^+} < +\infty.$$

From (29),

$$(61) \quad \begin{aligned} u(\rho\theta) \rho^{-\sigma_1^+} &= \rho^{-\sigma_1^+} \varphi_1(\rho) \psi_1(\theta) + \sum_{k=2}^{\bar{k}-1} \rho^{\sigma_k^+ - \sigma_1^+} \rho^{-\sigma_k^+} \varphi_k(\rho) \psi_k(\theta) \\ &+ \sum_{k=\bar{k}}^{\infty} \rho^{-\sigma_1^+} \varphi_k(\rho) \psi_k(\theta). \end{aligned}$$

From (51) we deduce that

$$(62) \quad \lim_{\rho \rightarrow 0^+} \rho^{-\sigma_1^+} \varphi_1(\rho) \psi_1(\theta) = \left[c_1^1 + \int_0^1 \frac{s^{-\sigma_1^+ + 1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds \right] \psi_1(\theta)$$

and

$$(63) \quad \lim_{\rho \rightarrow 0^+} \sum_{k=2}^{\bar{k}-1} \rho^{\sigma_k^+ - \sigma_1^+} \rho^{-\sigma_k^+} \varphi_k(\rho) \psi_k(\theta) = 0.$$

From (35), (58), (60), (59), and (57), we deduce that there exists some positive constant C_5 depending only on N , h , and a , such that, for all $\rho \in (0, 1/2)$,

$$\begin{aligned}
(64) \quad & \sum_{k=\bar{k}}^{\infty} \rho^{-\sigma_1^+} |\varphi_k(\rho)| |\psi_k(\theta)| \leq C_1 \sum_{k=\bar{k}}^{\infty} \rho^{-\sigma_1^+} \left(\frac{3}{2} |c_1^k| \rho^{\sigma_k^+} + |b_k| \rho^{\sigma_k^+/2} \right) |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \\
& \leq \frac{3}{2} C_1 \rho^{\sigma_2^+ - \sigma_1^+} \left(\sum_{k=\bar{k}}^{\infty} \frac{|c_1^k|^2}{N + 2\sigma_k^+} \right)^{\frac{1}{2}} \left(\sum_{k=\bar{k}}^{\infty} (N + 2\sigma_k^+) \rho^{2(\sigma_k^+ - \sigma_2^+)} |\mu_k|^2 \right)^{\frac{1}{2}} \\
& \quad + C_1 \rho^{(\sigma_k^+/2) - \sigma_1^+} \sum_{k=\bar{k}}^{\infty} |b_k| |\mu_k|^{\lfloor \frac{N-1}{4} \rfloor + 1} \\
& \leq C_5 \|u\|_{H^1(B(0,1))} (\rho^{2(\sigma_k^+ - \sigma_2^+)} + \rho^{(\sigma_k^+/2) - \sigma_1^+}),
\end{aligned}$$

which implies

$$(65) \quad \lim_{\rho \rightarrow 0^+} \sum_{k=\bar{k}}^{\infty} \rho^{-\sigma_1^+} \varphi_k(\rho) \psi_k(\theta) = 0.$$

Collecting (61), (62), (63), and (65), we finally obtain that

$$(66) \quad \lim_{\rho \rightarrow 0^+} u(\rho\theta) \rho^{-\sigma_1^+} = \left[c_1^1 + \int_0^1 \frac{s^{-\sigma_1^+ + 1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds \right] \psi_1(\theta).$$

We notice that, in view of (32), (30), and (40),

$$\begin{aligned}
(67) \quad & c_1^1 + \int_0^1 \frac{s^{-\sigma_1^+ + 1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds \\
& = \varphi_1(1) - c_2^1 + \int_0^1 \frac{s^{-\sigma_1^+ + 1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds \\
& = \int_{\mathbb{S}^{N-1}} u(\eta) \psi_1(\eta) dV(\eta) \\
& \quad + \int_{\mathbb{S}^{N-1}} \left[\int_0^1 \frac{s(s^{-\sigma_1^+} - s^{-\sigma_1^-})}{\sigma_1^+ - \sigma_1^-} h(s) u(s\eta) ds \right] \psi_1(\eta) dV(\eta).
\end{aligned}$$

The limit in (26) follows now from (66) and (67) in the case $r = 1$ and by a change of variable inside the integral for $r \neq 1$. Moreover estimates (44), (53), (55), (54), (56), the definition of j_k , and (64), imply that, for some $C_6 > 0$ depending only on N , R , h , and a ,

$$\begin{aligned}
(68) \quad & u(\rho\theta) \rho^{-\sigma_1^+} = \left(c_1^1 + \int_{\rho}^1 \frac{s^{-\sigma_k^+ + 1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_1(s) ds \right) \psi_1(\theta) \\
& \quad + \rho^{-\sigma_1^+} \left[B_1(\rho) \psi_1(\theta) + \sum_{k>1}^{\infty} \varphi_k(\rho) \psi_k(\theta) \right] \leq C_6 \|u\|_{H^1(B(0,R))},
\end{aligned}$$

for all $0 < \rho < \frac{1}{2}$. On the other hand, standard elliptic estimates in $B(0, 1) \setminus B(0, \frac{1}{2})$ yield, for some $C_7 > 0$ depending only on $N, R, h,$ and $a,$

$$(69) \quad u(\rho\theta)\rho^{-\sigma_1^+} \leq C_7 \|u\|_{H^1(B(0,R))}, \quad \text{for all } \frac{1}{2} \leq \rho < 1.$$

Estimate (28) follows from (68) and (69).

From (66) and the positivity of $u,$ it follows that $\left[c_1^1 + \int_0^1 \frac{s^{-\sigma_1^++1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds \right] \geq 0.$ Since

$$0 < \min_{\mathbb{S}^{N-1}} \psi_1 \leq \psi_1(\theta) \leq \max_{\mathbb{S}^{N-1}} \psi_1, \quad \text{for all } \theta \in \mathbb{S}^{N-1},$$

and, by standard regularity theory, $u \in C^0(\overline{B(0, 1)} \setminus B(0, s))$ for any $s \in (0, 1),$ the proof of Proposition 1 will be complete if we show that

$$(70) \quad c_1^1 + \int_0^1 \frac{s^{-\sigma_1^++1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds > 0.$$

In order to obtain (70), we need to prove the following

Claim 3: if $k \in \mathbb{N} \setminus \{0\}$ and

$$c_1^k + \int_0^1 \frac{s^{-\sigma_k^++1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds = 0.$$

then $\varphi_k(\rho) = 0$ for all $\rho \in (0, 1).$ Indeed, if $c_1^k + \int_0^1 \frac{s^{-\sigma_k^++1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds = 0,$ then

$$(71) \quad \varphi_k(\rho) = -\rho^{\sigma_k^+} \int_0^\rho \frac{s^{-\sigma_k^++1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds + \rho^{\sigma_k^-} \int_0^\rho \frac{s^{-\sigma_k^-+1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds.$$

From claim 2 and (51), we know that there exists a constant ℓ_k depending on k (and on a, R, h, u, N) such that

$$|\varphi_k(\rho)| \leq \ell_k \rho^{\sigma_k^+}.$$

Using the above estimate in (71), we can improve such an estimate as

$$|\varphi_k(\rho)| \leq \ell_k \rho^{\sigma_k^+} \frac{2 \|h\|_{L^p(B(0,1))} \rho^\varepsilon}{\omega_N (\sigma_k^+ - \sigma_k^-) \varepsilon}.$$

Using the above estimate in (71), we can obtain the following further improvement

$$|\varphi_k(\rho)| \leq \ell_k \rho^{\sigma_k^+} \left(\frac{2 \|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-) \varepsilon} \right)^2 \frac{(\rho^\varepsilon)^2}{(2 \cdot 1)^{\frac{p-1}{p}}}.$$

Arguing by induction, we can easily prove that, for all $j \in \mathbb{N},$

$$|\varphi_k(\rho)| \leq \ell_k \rho^{\sigma_k^+} \left(\frac{2 \|h\|_{L^p(B(0,1))}}{\omega_N (\sigma_k^+ - \sigma_k^-) \varepsilon} \right)^j \frac{(\rho^\varepsilon)^j}{(j!)^{\frac{p-1}{p}}},$$

and letting $j \rightarrow +\infty,$ we deduce that $\varphi_k(\rho) = 0$ for all $\rho \in (0, 1).$ Claim 3 is thereby proved.

We are now in position to prove (70). Arguing by contradiction, let us assume that

$$(72) \quad c_1^1 + \int_0^1 \frac{s^{-\sigma_1^++1}}{\sigma_1^+ - \sigma_1^-} h(s) \varphi_1(s) ds = 0$$

and let $k_0 > 1$ be the smallest index for which

$$c_1^{k_0} + \int_0^1 \frac{s^{-\sigma_{k_0}^+ + 1}}{\sigma_{k_0}^+ - \sigma_{k_0}^-} h(s) \varphi_{k_0}(s) ds \neq 0.$$

Such a k_0 exists in view of claim 3; indeed if $c_1^k + \int_0^1 \frac{s^{-\sigma_k^+ + 1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds = 0$ for all k , then $\varphi_k \equiv 0$ for all k and u would be identically zero, thus giving rise to a contradiction. Moreover, from (72), we have that $k_0 > 1$, and, by claim 3, $\varphi_k \equiv 0$ in $(0, 1)$ for all $1 \leq k \leq k_0 - 1$. Repeating the same arguments we used above to prove (66), it is now possible to show that

$$(73) \quad \lim_{\rho \rightarrow 0^+} u(\rho\theta) \rho^{-\sigma_{k_0}^+} = \sum_{k=k_0}^{k_0+m_{k_0}-1} \left[c_1^k + \int_0^1 \frac{s^{-\sigma_k^+ + 1}}{\sigma_k^+ - \sigma_k^-} h(s) \varphi_k(s) ds \right] \psi_k(\theta),$$

where m_{k_0} is the geometric multiplicity of the eigenvalue μ_{k_0} . We notice that the sum at the right hand side is a nontrivial function in $L^2(\mathbb{S}^{N-1})$ which, being $k_0 > 1$, is orthogonal to the first positive eigenfunction ψ_1 . Hence the right hand side of (73) changes sign in \mathbb{S}^{N-1} . Therefore the limit in (73) implies that u changes sign in a neighborhood of 0, which is in contradiction with the positivity assumption on u . Condition (70) follows and the proof of Proposition 1 is now complete. \square

Remark 2. If we let the assumption of positivity of u drop, following the proof of Proposition 1, we can still prove a *Cauchy's integral type formula* for u . More precisely, if $u \in H^1(B(0, R))$ is a weak solution to (25) in $B(0, R)$ which changes sign in any neighborhood of 0, with a radial potential $h \in L_{\text{loc}}^\infty(0, R) \cap L^p(0, R)$ for some $p > N/2$, then, following the notation introduced in (33) and letting $k_0 > 1$ be the smallest index for which

$$\int_{\mathbb{S}^{N-1}} \left(r^{-\sigma_{k_0}^+} u(r\eta) + \int_0^r \frac{s^{1-\sigma_{k_0}^+}}{\sigma_{k_0}^+ - \sigma_{k_0}^-} h(s) u(s\eta) ds - r^{\sigma_{k_0}^- - \sigma_{k_0}^+} \int_0^r \frac{s^{1-\sigma_{k_0}^-}}{\sigma_{k_0}^+ - \sigma_{k_0}^-} h(s) u(s\eta) ds \right) \psi_{k_0}(\eta) dV(\eta) \neq 0,$$

for any $\theta \in \mathbb{S}^{N-1}$ and $r \in (0, R)$ there holds

$$\lim_{\rho \rightarrow 0^+} u(\rho\theta) \rho^{-\sigma_{k_0}^+} = \sum_{\{k: \mu_k = \mu_{k_0}\}} \left[\int_{\mathbb{S}^{N-1}} \left(r^{-\sigma_k^+} u(r\eta) + \int_0^r \frac{s^{1-\sigma_k^+}}{\sigma_k^+ - \sigma_k^-} h(s) u(s\eta) ds - r^{\sigma_k^- - \sigma_k^+} \int_0^r \frac{s^{1-\sigma_k^-}}{\sigma_k^+ - \sigma_k^-} h(s) u(s\eta) ds \right) \psi_k(\eta) dV(\eta) \right] \psi_k(\theta).$$

Without the assumption of radial symmetry of the potential, it is still possible to evaluate the exact behavior near the singularity of the first Fourier coefficient φ_1 (see (29) and (30)).

Lemma 4.1. *Let $a \in L^\infty(\mathbb{S}^{N-1})$ be such that $\Lambda_N(a) < 1$, $R > 0$, and $u \geq 0$ a.e. in $B(0, R)$, $u \in H^1(B(0, R))$, $u \not\equiv 0$, be a weak H^1 -solution to*

$$-\Delta u(x) = \left[\frac{a(x/|x|)}{|x|^2} + q(x) \right] u(x) \quad \text{in } B(0, R),$$

where $q \in L_{\text{loc}}^\infty(B(0, R) \setminus \{0\}) \cap L^p(B(0, R))$ for some $p > \frac{N}{2}$. Then, for any $0 < r < R$,

$$(74) \quad \begin{aligned} & \lim_{\rho \rightarrow 0^+} \rho^{-\sigma} \int_{\mathbb{S}^{N-1}} u(\rho\theta)\psi_1(\theta) dV(\theta) \\ &= \int_{\mathbb{S}^{N-1}} \left(r^{-\sigma} u(r\theta) + \int_0^r \frac{s^{1-\sigma}}{2\sigma + N - 2} q(s\theta)u(s\theta) ds \right. \\ & \quad \left. - r^{-2\sigma - N + 2} \int_0^r \frac{s^{N-1+\sigma}}{2\sigma + N - 2} q(s\theta)u(s\theta) ds \right) \psi_1(\theta) dV(\theta). \end{aligned}$$

Proof. The proof follows the lines of the first part of the proof of Proposition 1. By scaling, it is sufficient to prove (74) for $r = 1$. Let

$$u(x) = u(\rho\theta) = \sum_{k=1}^{\infty} \varphi_k(\rho)\psi_k(\theta) \quad \text{and} \quad q(x)u(x) = q(\rho\theta)u(\rho\theta) = \sum_{k=1}^{\infty} \zeta_k(\rho)\psi_k(\theta)$$

where $\rho = |x| \in (0, 1]$, $\theta = x/|x| \in \mathbb{S}^{N-1}$,

$$\varphi_k(\rho) = \int_{\mathbb{S}^{N-1}} u(\rho\theta)\psi_k(\theta) dV(\theta), \quad \zeta_k(\rho) = \int_{\mathbb{S}^{N-1}} q(\rho\theta)u(\rho\theta)\psi_k(\theta) dV(\theta),$$

and ψ_k is an L^2 -normalized eigenfunction of the operator $-\Delta_{\mathbb{S}^{N-1}} - a(\theta)$ on the sphere associated to the k -th eigenvalue μ_k , i.e. satisfying (11). The first Fourier coefficient φ_1 solves

$$\varphi_1''(\rho) + \frac{N-1}{\rho} \varphi_1'(\rho) - \frac{\mu_1}{\rho^2} \varphi_1(\rho) = \zeta_1(\rho) \quad \text{in } (0, 1).$$

A direct calculation shows that, for some $c_1^+, c_2^+ \in \mathbb{R}$,

$$\varphi_1(\rho) = \rho^{\sigma_1^+} \left(c_1^+ + \int_\rho^1 \frac{s^{-\sigma_1^+ + 1}}{\sigma_1^+ - \sigma_1^-} \zeta_1(s) ds \right) + \rho^{\sigma_1^-} \left(c_2^+ + \int_\rho^1 \frac{s^{-\sigma_1^- + 1}}{\sigma_1^- - \sigma_1^+} \zeta_1(s) ds \right),$$

where $\sigma_1^+ = -\frac{N-2}{2} + \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1}$ and $\sigma_1^- = -\frac{N-2}{2} - \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1}$. From Theorem 1.2 and standard regularity theory, we deduce that $u(x) \leq \text{const } |x|^{\sigma_1^+}$ in $B(0, 1)$, hence Hölder's inequality yields

$$(75) \quad \begin{aligned} & \int_0^1 |s^{-\sigma_1^+ + 1} \zeta_1(s)| ds \leq \int_0^1 s^{-\sigma_1^+ + 1} \left(\int_{\mathbb{S}^{N-1}} |q(s\theta)|u(s\theta)\psi_1(\theta) dV(\theta) \right) ds \\ & \leq \text{const} \int_0^1 s^{1 - \frac{N-1}{p}} \left(\int_{\mathbb{S}^{N-1}} s^{\frac{N-1}{p}} |q(s\theta)| dV(\theta) \right) ds \\ & \leq \text{const} \|q\|_{L^p(B(0,1))} \left(\int_0^1 s^{(1 - \frac{N-1}{p}) \frac{p}{p-1}} ds \right)^{1 - \frac{1}{p}} < \infty. \end{aligned}$$

In a similar way, we obtain that

$$(76) \quad s \mapsto \frac{s^{-\sigma_1^- + 1}}{\sigma_1^- - \sigma_1^+} \zeta_1(s) \in L^1(0, 1).$$

Since $u \in L^{2^*}(B(0, 1))$, $\sigma_1^- < \sigma_1^+$, from (75), (76), and the fact that $\rho^{\sigma_1^-} \psi_1(\theta)$ does not belong to $L^{2^*}(B(0, 1))$, we conclude that there must be

$$c_2^1 = - \int_0^1 \frac{s^{-\sigma_1^-+1}}{\sigma_1^- - \sigma_1^+} \zeta_1(s) ds,$$

hence

$$(77) \quad \rho^{-\sigma_1^+} \varphi_1(\rho) = c_1^1 + \int_\rho^1 \frac{s^{-\sigma_1^++1}}{\sigma_1^+ - \sigma_1^-} \zeta_1(s) ds + \rho^{\sigma_1^- - \sigma_1^+} \int_0^\rho \frac{s^{-\sigma_1^-+1}}{\sigma_1^+ - \sigma_1^-} \zeta_1(s) ds$$

for any $\rho \in (0, 1)$. Notice that, by standard regularity theory, φ_1 is continuous at $\rho = 1$, thus, letting $\rho \rightarrow 1^-$ in (77), we obtain

$$(78) \quad c_1^1 = \varphi_1(1) - \int_0^1 \frac{s^{-\sigma_1^-+1}}{\sigma_1^+ - \sigma_1^-} \zeta_1(s) ds.$$

Arguing as in (75), we obtain that

$$(79) \quad \rho^{\sigma_1^- - \sigma_1^+} \int_0^\rho \left| \frac{s^{-\sigma_1^-+1}}{\sigma_1^- - \sigma_1^+} \zeta_1(s) \right| ds \leq \text{const } \rho^{2 - \frac{N}{p}}.$$

Since $p > \frac{N}{2}$, (77), (78), and (79) imply that

$$\lim_{\rho \rightarrow 0^+} \rho^{-\sigma_1^+} \varphi_1(\rho) = \varphi_1(1) - \int_0^1 \frac{s^{-\sigma_1^-+1}}{\sigma_1^+ - \sigma_1^-} \zeta_1(s) ds + \int_0^1 \frac{s^{-\sigma_1^++1}}{\sigma_1^+ - \sigma_1^-} \zeta_1(s) ds,$$

and (74) for $r = 1$ follows. The result in the case $r \neq 1$ can be easily obtained just by scaling. \square

In order to extend the result of Proposition 1 to the case in which the potential is a non radial perturbation of the dipole-type singular term, we will construct a subsolution and a supersolution which solve equations of type (25) and the behavior of which is consequently known in view of Proposition 1.

Lemma 4.2. *Let $a \in L^\infty(\mathbb{S}^{N-1})$ be such that $\Lambda_N(a) < 1$, $C \in \mathbb{R}$, and $\varepsilon > 0$. Then, for all*

$$(80) \quad 0 < r < \begin{cases} \left[\frac{(N-2)^2}{4C^+} (1 - \Lambda_N(a)) \right]^{1/\varepsilon}, & \text{if } C > 0, \\ +\infty, & \text{if } C \leq 0, \end{cases}$$

and for all $\gamma \in H^{1/2}(\partial B(0, r))$, $\gamma \geq 0$, $\gamma \not\equiv 0$, the Dirichlet boundary value problem

$$(81) \quad \begin{cases} -\Delta u(x) = \left[\frac{a(x/|x|)}{|x|^2} + C|x|^{-2+\varepsilon} \right] u(x), & \text{in } B(0, r), \\ u|_{\partial B(0, r)} = \gamma, & \text{on } \partial B(0, r), \end{cases}$$

admits a unique weak solution $u \in H^1(B(0, r))$. Moreover u is continuous and strictly positive in $B(0, r) \setminus \{0\}$, and there exists a positive constant C' depending on a , C , ε , N , and r , such that

$$(82) \quad \|u\|_{H^1(B(0, r))} \leq C' \|\gamma\|_{H^{1/2}(\partial B(0, r))}.$$

In addition, if $\gamma \in W^{2-\frac{1}{k}, k}(\partial B(0, r))$ for some $k > N/2$, then $u \in C^0(\overline{B(0, r)} \setminus \{0\})$.

Proof. For a fixed r satisfying (80) and $\gamma \in H^{1/2}(\partial B(0, r))$, $\gamma \geq 0$, $\gamma \not\equiv 0$, let \tilde{v} be the unique $H^1(B(0, r))$ -weak solution to the problem

$$\begin{cases} \tilde{v} \in H^1(B(0, r)), \\ -\Delta \tilde{v} = 0, & \text{in } B(0, r), \\ v = \gamma, & \text{on } \partial B(0, r). \end{cases}$$

By classical trace embedding theorems, it follows that

$$(83) \quad \|\tilde{v}\|_{H^1(B(0, r))} \leq \text{const}(N, r) \|\gamma\|_{H^{1/2}(\partial B(0, r))},$$

for some positive constant $\text{const}(N, r)$ depending only on N and r . Let us define the quadratic form $a : H_0^1(B(0, r)) \times H_0^1(B(0, r)) \rightarrow \mathbb{R}$ as

$$\mathcal{Q}(w, u) := \int_{B(0, r)} \left[\nabla w(x) \cdot \nabla u(x) - \frac{1}{|x|^2} (a(x/|x|) + C|x|^\varepsilon) w(x)u(x) \right] dx,$$

and $\Phi \in H^{-1}(B(0, r))$ as

$$H^{-1}\langle \Phi, u \rangle_{H_0^1} := \int_{B(0, r)} \left(\frac{a(x/|x|)}{|x|^2} + \frac{C}{|x|^{2-\varepsilon}} \right) \tilde{v}(x)u(x) dx.$$

By Hardy's inequality, it is easy to verify that

$$(84) \quad \mathcal{Q}(u, u) \geq \left[1 - \Lambda_N(a) - \frac{4C^+r^\varepsilon}{(N-2)^2} \right] \int_{B(0, r)} |\nabla u(x)|^2 dx.$$

Since (80) implies that $\left[1 - \Lambda_N(a) - \frac{4C^+r^\varepsilon}{(N-2)^2} \right] > 0$, we conclude that the bilinear bounded form \mathcal{Q} is coercive. Furthermore, the function $x \mapsto a\left(\frac{x}{|x|}\right)|x|^{-2} + C|x|^{-2+\varepsilon}$ belongs to $L^{\frac{2N}{N+2}}(B(0, r))$, hence Φ is a bounded linear functional on $H_0^1(B(0, r))$. From the Lax-Milgram lemma we deduce that there exists a unique $w \in H_0^1(B(0, r))$ such that $\mathcal{Q}(w, u) = H^{-1}\langle \Phi, u \rangle_{H_0^1}$ for all $u \in H_0^1(B(0, r))$. In particular w weakly solves

$$(85) \quad \begin{cases} -\Delta w(x) - \frac{1}{|x|^2} \left[a\left(\frac{x}{|x|}\right) + C|x|^\varepsilon \right] w(x) = \left[\frac{a(x/|x|)}{|x|^2} + \frac{C}{|x|^{2-\varepsilon}} \right] \tilde{v}(x), & \text{in } B(0, r), \\ w = 0, & \text{on } \partial B(0, r). \end{cases}$$

Testing the above equation with w and using (84), Poincaré's and Hölder's inequalities and (83), we obtain that

$$\|w\|_{H^1(B(0, r))} \leq c(a, C, \varepsilon, N, r) \|\tilde{v}\|_{H^1(B(0, r))} \leq c'(a, C, \varepsilon, N, r) \|\gamma\|_{H^{1/2}(\partial B(0, r))},$$

for some positive constants $c(a, C, \varepsilon, N, r)$ and $c'(a, C, \varepsilon, N, r)$ depending on a, C, ε, N , and r . It is now easy to verify that $u := w + \tilde{v} \in H^1(B(0, r))$ satisfies (82) and is the unique weak solution to (81). Moreover, testing (81) with $-u^- := -\max\{-u, 0\}$ and using (84), we obtain that

$$0 = \mathcal{Q}(u^-, u^-) \geq \left[1 - \Lambda_N(a) - \frac{4C^+r^\varepsilon}{(N-2)^2} \right] \int_{B(0, r)} |\nabla u^-(x)|^2 dx,$$

which, in view of (80), implies that $u^- = 0$ a.e. in $B(0, r)$, i.e. $u \geq 0$ a.e. in $B(0, r)$. The Strong Maximum Principle allows us to conclude that $u > 0$ in $B(0, r) \setminus \{0\}$, while standard regularity theory for elliptic equations ensures interior continuity of u outside the origin.

If, in addition, we assume that $\gamma \in W^{2-1/k,k}(\partial B(0,r))$ for some $k > N/2$, then $\tilde{v} \in W^{2,k}(B(0,r))$, and hence $\tilde{v} \in C^{0,\alpha}(\overline{B(0,r)})$, hence, from elliptic regularity theory applied to (85) outside 0, we obtain that $u \in C^0(\overline{B(0,r)} \setminus \{0\})$. \square

Proof of Theorem 1.1. Let $R > 0$ such that $\overline{B(0,R)} \subset \Omega$. Since $q(x) = O(|x|^{-(2-\varepsilon)})$ as $|x| \rightarrow 0$ for some $\varepsilon > 0$ and $q \in L_{\text{loc}}^\infty(\Omega \setminus \{0\})$, there exists a positive constant \tilde{C} such that $-\tilde{C}|x|^{-(2-\varepsilon)} \leq q(x) \leq \tilde{C}|x|^{-(2-\varepsilon)}$ for a.e. $x \in B(0,R)$. Let us fix $\bar{r} = \bar{r}(R, N, q, a, \varepsilon)$, such that $0 < \bar{r} < \min \left\{ R, \left[\frac{(N-2)^2}{4\tilde{C}} (1 - \Lambda_N(a)) \right]^{1/\varepsilon} \right\}$. We notice that the Maximum Principle implies that $u > 0$ in $\overline{B(0,R)} \setminus \{0\}$, whereas standard elliptic regularity theory yields $u \in W^{2,k}(\overline{B(0,R)} \setminus B(0,s))$ for all $s \in (0,R)$ and some $k > N/2$, and, consequently, u is continuous in $\overline{B(0,R)} \setminus \{0\}$. Hence the function $\gamma_r := u|_{\partial B(0,r)}$ belongs to $W^{2-1/k,k}(\partial B(0,r))$ for some $k > N/2$ and is continuous and strictly positive on $\partial B(0,r)$ for all $0 < r \leq \bar{r}$. From Lemma 4.2 we deduce that, for any $0 < r \leq \bar{r}$, there exist $\underline{u}_r \in H^1(B(0,r))$ and $\bar{u}_r \in H^1(B(0,r))$ continuous and strictly positive in $\overline{B(0,r)} \setminus \{0\}$, weakly satisfying

$$\begin{cases} -\Delta \underline{u}_r(x) = \left[\frac{a(x/|x|)}{|x|^2} - \tilde{C}|x|^{-2+\varepsilon} \right] \underline{u}_r(x), & \text{in } B(0,r), \\ \underline{u}_r|_{\partial B(0,r)} = \gamma_r, & \text{on } \partial B(0,r), \end{cases}$$

and

$$\begin{cases} -\Delta \bar{u}_r(x) = \left[\frac{a(x/|x|)}{|x|^2} + \tilde{C}|x|^{-2+\varepsilon} \right] \bar{u}_r(x), & \text{in } B(0,r), \\ \bar{u}_r|_{\partial B(0,r)} = \gamma_r, & \text{on } \partial B(0,r). \end{cases}$$

From Proposition 1, there exist two constants $A_2 > A_1 > 0$ (depending on $\bar{r}, N, q, a, \varepsilon$, and u) such that

$$(86) \quad A_1|x|^\sigma \leq \underline{u}_{\bar{r}}(x) \quad \text{and} \quad \bar{u}_{\bar{r}}(x) \leq A_2|x|^\sigma, \quad \text{for all } x \in B(0, \bar{r}/2) \setminus \{0\}.$$

Furthermore, for all $0 < r \leq \bar{r}$, $u - \underline{u}_r$ satisfies

$$(87) \quad \begin{cases} -\Delta(u - \underline{u}_r)(x) - \left[\frac{a(x/|x|)}{|x|^2} - \tilde{C}|x|^{-2+\varepsilon} \right] (u - \underline{u}_r)(x) \geq 0, & \text{in } B(0,r), \\ (u - \underline{u}_r)|_{\partial B(0,r)} = 0, & \text{on } \partial B(0,r), \end{cases}$$

while $u - \bar{u}_r$ satisfies

$$(88) \quad \begin{cases} -\Delta(u - \bar{u}_r)(x) - \left[\frac{a(x/|x|)}{|x|^2} + \tilde{C}|x|^{-2+\varepsilon} \right] (u - \bar{u}_r)(x) \leq 0, & \text{in } B(0,r), \\ (u - \bar{u}_r)|_{\partial B(0,r)} = 0, & \text{on } \partial B(0,r). \end{cases}$$

Testing (87), respectively (88), with $-(u - \underline{u}_r)^-$, respectively $(u - \bar{u}_r)^+$, and using (84), we obtain that, for any $0 < r \leq \bar{r}$,

$$(89) \quad \underline{u}_r(x) \leq u(x) \leq \bar{u}_r(x), \quad \text{for all } x \in B(0,r) \setminus \{0\}.$$

In particular, from (86) and (89), we deduce that

$$(90) \quad A_1|x|^\sigma \leq u(x) \leq A_2|x|^\sigma, \quad \text{for all } x \in B(0, \bar{r}/2) \setminus \{0\}.$$

Estimate (90) and the continuity of u outside the origin imply that there exists a positive constant C (depending on $q, R, \Omega, a, \varepsilon$, and u) such that

$$(91) \quad \frac{1}{C}|x|^\sigma \leq u(x) \leq C|x|^\sigma \quad \text{for all } x \in \overline{B(0, R)} \setminus \{0\}.$$

Let us now fix $\delta = \delta(N, a, \varepsilon) > 0$ such that

$$\delta < \min \left\{ \varepsilon, \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1} \right\}$$

and set

$$\hat{r} = \min \left\{ \bar{r}, \left[\frac{\delta}{C} \left(2\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1} - \delta \right) \right]^{1/\varepsilon}, \left(\frac{\min_{\mathbb{S}^{N-1}} \psi_1}{C} \right)^{1/\delta} \right\},$$

with C given in (91). The function \hat{u} defined as

$$(92) \quad \hat{u}(x) = |x|^{\sigma-\delta} \psi_1(x/|x|)$$

belongs to $H^1(B(0, R))$ and, for all $0 < r \leq \hat{r}$, satisfies

$$\begin{cases} -\Delta \hat{u}(x) - \frac{a(x/|x|)}{|x|^2} \hat{u}(x) = \delta \left(2\sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_1} - \delta \right) |x|^{-2} \hat{u}(x) \\ \qquad \qquad \qquad \geq \tilde{C} |x|^{-2+\varepsilon} \hat{u}(x), & \text{in } B(0, r), \\ \hat{u}(x)|_{\partial B(0, r)} = r^{\sigma-\delta} \psi_1(x/r) \geq \bar{u}_r(x), & \text{on } \partial B(0, r). \end{cases}$$

Hence, for all $0 < r \leq \hat{r}$, $\hat{u} - \bar{u}_r$ satisfies

$$\begin{cases} -\Delta(\hat{u} - \bar{u}_r)(x) - \left[\frac{a(x/|x|)}{|x|^2} x \cdot \mathbf{d} + \tilde{C} |x|^{-2+\varepsilon} \right] (\hat{u} - \bar{u}_r)(x) \geq 0, & \text{in } B(0, r), \\ (\hat{u} - \bar{u}_r)|_{\partial B(0, r)} \geq 0, & \text{on } \partial B(0, r). \end{cases}$$

Testing the above equation with $-(\hat{u} - \bar{u}_r)^-$ and using (84), we obtain that, for any $0 < r \leq \hat{r}$,

$$(93) \quad \bar{u}_r(x) \leq \hat{u}(x), \quad \text{for all } x \in B(0, r) \setminus \{0\}.$$

From Proposition 1, for any $0 < r \leq \hat{r}$, the functions

$$x \mapsto \frac{\underline{u}_r(x)}{|x|^\sigma \psi_1(x/|x|)} \quad \text{and} \quad x \mapsto \frac{\bar{u}_r(x)}{|x|^\sigma \psi_1(x/|x|)}$$

have limits as $|x| \rightarrow 0$, which, accordingly with (26–27) and taking into account the continuity of functions \underline{u}_r and \bar{u}_r up to the boundary $|x| = r$, can be computed as

$$\begin{aligned} \underline{L}_r &:= \lim_{|x| \rightarrow 0} \frac{\underline{u}_r(x)}{|x|^\sigma \psi_1(x/|x|)} \\ &= \int_{\mathbb{S}^{N-1}} \left(r^{-\sigma} u(r\eta) - \tilde{C} \int_0^r \frac{s^{1-\sigma}}{2\sigma + N - 2} s^{-2+\varepsilon} \underline{u}_r(s\eta) ds \right. \\ &\quad \left. + \tilde{C} r^{-2\sigma - N + 2} \int_0^r \frac{s^{N-1+\sigma}}{2\sigma + N - 2} s^{-2+\varepsilon} \underline{u}_r(s\eta) ds \right) \psi_1(\eta) dV(\eta), \end{aligned}$$

and

$$\begin{aligned}\bar{L}_r &:= \lim_{|x| \rightarrow 0} \frac{\bar{u}_r(x)}{|x|^\sigma \psi_1(x/|x|)} \\ &= \int_{\mathbb{S}^{N-1}} \left(r^{-\sigma} u(r\eta) + \tilde{C} \int_0^r \frac{s^{1-\sigma}}{2\sigma + N - 2} s^{-2+\varepsilon} \bar{u}_r(s\eta) ds \right. \\ &\quad \left. - \tilde{C} r^{-2\sigma-N+2} \int_0^r \frac{s^{N-1+\sigma}}{2\sigma + N - 2} s^{-2+\varepsilon} \bar{u}_r(s\eta) ds \right) \psi_1(\eta) dV(\eta).\end{aligned}$$

From (89) and (91), it follows that

$$\underline{L}_r = r^{-\sigma} \int_{\mathbb{S}^{N-1}} u(r\eta) \psi_1(\eta) dV(\eta) + o(1) \quad \text{as } r \rightarrow 0.$$

From (93), (92), and the choice of δ , we obtain that

$$\bar{L}_r = r^{-\sigma} \int_{\mathbb{S}^{N-1}} u(r\eta) \psi_1(\eta) dV(\eta) + o(1) \quad \text{as } r \rightarrow 0.$$

Hence, from Lemma 4.1, we conclude that, for any R such that $\overline{B(0, R)} \subset \Omega$,

$$(94) \quad \begin{aligned}\lim_{r \rightarrow 0} \underline{L}_r &= \lim_{r \rightarrow 0} \bar{L}_r = \int_{\mathbb{S}^{N-1}} \left(R^{-\sigma} u(R\eta) + \int_0^R \frac{s^{1-\sigma}}{2\sigma + N - 2} q(s\eta) u(s\eta) ds \right. \\ &\quad \left. - R^{-2\sigma-N+2} \int_0^R \frac{s^{N-1+\sigma}}{2\sigma + N - 2} q(s\eta) u(s\eta) ds \right) \psi_1(\eta) dV(\eta).\end{aligned}$$

In view of (89), there holds that, for any $0 < r \leq \bar{r}$,

$$\begin{aligned}\underline{L}_r &= \lim_{|x| \rightarrow 0} \frac{\underline{u}_r(x)}{|x|^{a_{\mu_1}} \psi_1(x/|x|)} \leq \liminf_{|x| \rightarrow 0} \frac{u(x)}{|x|^{a_{\mu_1}} \psi_1(x/|x|)} \\ &\leq \limsup_{|x| \rightarrow 0} \frac{u(x)}{|x|^{a_{\mu_1}} \psi_1(x/|x|)} \leq \lim_{|x| \rightarrow 0} \frac{\bar{u}_r(x)}{|x|^{a_{\mu_1}} \psi_1(x/|x|)} = \bar{L}_r.\end{aligned}$$

Letting $r \rightarrow 0$, we complete the proof. \square

5. BEHAVIOR OF SOLUTIONS TO THE SEMILINEAR PROBLEM

The L^q and L^∞ bounds of solutions to dipole-type linear Schrödinger equations with properly summable potentials, derived in Theorems 1.2 and 3.2, allow us to obtain in the semilinear case analogous estimates.

Theorem 5.1. *Let Ω be a bounded domain containing 0, $a \in L^\infty(\mathbb{S}^{N-1})$ such that $\Lambda_N(a) < 1$, and $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ such that, for some positive constant C ,*

$$\left| \frac{f(x, u)}{u} \right| \leq C (1 + |u|^{2^*-2}) \quad \text{for a.e. } (x, u) \in \Omega \times \mathbb{R}.$$

Then, for any $\Omega' \Subset \Omega$ and for any weak $H^1(\Omega)$ -solution u of (7), there holds $\frac{u}{\varphi} \in L^\infty(\Omega')$.

Proof. Let $\Omega' \Subset \Omega$ and $u \in H^1(\Omega)$ be a weak $H^1(\Omega)$ -solution to (7). We set

$$V(x) = \frac{f(x, u(x))}{\varphi^{2^*-2}(x)u(x)}$$

and notice that $u \in L^{2^*}(\Omega)$ yields

$$\int_{\Omega} \varphi^{2^*}(x) |V(x)|^{N/2} dx < +\infty.$$

Hence Theorem 3.2 implies that $\frac{u}{\varphi} \in L^q(\varphi^{2^*}, \Omega')$ for all $1 \leq q < +\infty$. Since

$$\int_{\Omega} \varphi^{2^*}(x) |V(x)|^s dx \leq \text{const} \left(1 + \int_{\Omega} \varphi^{2^*}(x) \left| \frac{u(x)}{\varphi(x)} \right|^{(2^*-2)s} dx \right),$$

we obtain that $V \in L^s(\varphi^{2^*}, \Omega)$ for all $s \geq \frac{N-2}{4}$. The conclusion follows now from Theorem 1.2. \square

Proof of Theorem 1.3. From Theorem 5.1, it follows that $q(x) = \frac{f(x, u(x))}{u(x)}$ is such that $q \in L_{\text{loc}}^{\infty}(\Omega \setminus \{0\})$ and $q(x) = O(|x|^{-(2-\varepsilon)})$ as $|x| \rightarrow 0$ for some $\varepsilon > 0$. Hence the conclusion follows from Theorem 1.1. \square

REFERENCES

- [1] B. Abdellaoui, V. Felli and I. Peral, *Existence and multiplicity for perturbations of an equation involving Hardy inequality and critical Sobolev exponent in the whole \mathbb{R}^N* , Adv. Differential Equations, **9** (2004), 481–508.
- [2] H. Brezis and T. Kato, *Remarks on the Schrödinger operator with singular complex potentials*, J. Math. Pures Appl. (9), **58** (1979), 137–151.
- [3] L. Caffarelli, R. Kohn and L. Nirenberg, *First order interpolation inequalities with weights*, Compositio Math., **53** (1984), 259–275.
- [4] F. Catrina and Z.-Q. Wang, *On the Caffarelli-Kohn-Nirenberg inequalities: sharp constants, existence (and nonexistence), and symmetry of extremal functions*, Comm. Pure Appl. Math., **54** (2001), 229–258.
- [5] V. De Cicco and M. A. Vivaldi, *A Liouville type theorem for weighted elliptic equations*, Adv. Math. Sci. Appl., **9** (1999), 183–207.
- [6] H. Egnell, *Elliptic boundary value problems with singular coefficients and critical nonlinearities*, Indiana Univ. Math. J., **38** (1989), 235–251.
- [7] V. Felli, E.M. Marchini and S. Terracini, *On Schrödinger operators with multipolar inverse-square potentials*, J. Funct. Anal., to appear.
- [8] V. Felli and M. Schneider, *A note on regularity of solutions to degenerate elliptic equations of Caffarelli-Kohn-Nirenberg type*, Adv. Nonlinear Stud., **3** (2003), 431–443.
- [9] J. García Azorero and I. Peral, *Hardy inequalities and some critical elliptic and parabolic problems*, J. Diff. Equations, **144** (1998), 441–476.
- [10] E. Jannelli, *The role played by space dimension in elliptic critical problems*, J. Differential Equations, **156** (1999), 407–426.
- [11] J. M. Lévy-Leblond, *Electron capture by polar molecules*, Phys. Rev., **153** (1967), 1–4.
- [12] M. Murata, *Structure of positive solutions to $(-\Delta + V)u = 0$ in \mathbb{R}^n* , Duke Math. J., **53** (1986), 869–943.
- [13] Y. Pinchover, *On positive Liouville theorems and asymptotic behavior of solutions of Fuchsian type elliptic operators*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **11** (1994), 313–341.
- [14] M. Reed and B. Simon, “Methods of Modern Mathematical Physics. IV. Analysis of Operators,” Academic Press, New York-London, 1978.
- [15] Yu. Safarov and D. Vassiliev, “The Asymptotic Distribution of Eigenvalues of Partial Differential Operators,” Translations of Mathematical Monographs, 155, American Mathematical Society, Providence, RI, 1997.
- [16] B. Simon, *Schrödinger operators in the twentieth century*, J. Math. Phys., **41** (2000), 3523–3555.
- [17] D. Smets, *Nonlinear Schrödinger equations with Hardy potential and critical nonlinearities*, Trans. AMS, **357** (2005), 2909–2938.
- [18] S. Terracini, *On positive entire solutions to a class of equations with singular coefficient and critical exponent*, Adv. Differential Equations, **1** (1996), 241–264.

Received December 2006; revised April 2007.

VERONICA FELLI: UNIVERSITÀ DI MILANO BICOCCA, DIPARTIMENTO DI STATISTICA, VIA BICOCCA DEGLI ARCIMBOLDI 8, 20126 MILANO, ITALY. *E-mail address:* `veronica.felli@unimib.it`.

ELSA M. MARCHINI, SUSANNA TERRACINI: UNIVERSITÀ DI MILANO BICOCCA, DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, VIA COZZI 53, 20125 MILANO, ITALY. *E-mail addresses:* `elsa.marchini@unimib.it`, `susanna.terracini@unimib.it`.