

# Nodal domains and spectral minimal partitions.

B. HELFFER<sup>1,3</sup>

T. HOFFMANN-OSTENHOF<sup>2,3</sup>

S. TERRACINI<sup>4</sup>

Département de Mathématiques, UMR 8628 Univ Paris-Sud<sup>1</sup>

Institut für Theoretische Chemie, Universität Wien<sup>2</sup>

International Erwin Schrödinger Institute for Mathematical Physics<sup>3</sup>

Università di Milano Bicocca, Dipartimento di Matematica e Applicazioni<sup>4</sup>

October 23, 2006

## Abstract

We consider two-dimensional Schrödinger operators in bounded domains. We analyze relations between the nodal domains, spectral minimal partitions and spectral properties of the corresponding operator. The main results concern the existence and regularity of the minimal partitions and the characterization of the minimal partitions associated with nodal sets as the nodal domains of Courant–sharp eigenfunctions.

# Contents

<b>1</b>	<b>Introduction and main results</b>	<b>3</b>
1.1	Definitions and notations . . . . .	3
1.2	Main results . . . . .	6
1.3	Organization of the paper . . . . .	9
<b>2</b>	<b>Preliminaries: Hölder regularity of nodal sets</b>	<b>9</b>
<b>3</b>	<b>Optimal partitions in <math>N</math> dimensions</b>	<b>16</b>
<b>4</b>	<b>Further results in two dimensions</b>	<b>25</b>
4.1	Case when $\mathbf{k}_0 = \emptyset$ . . . . .	26
4.2	General case. . . . .	35
<b>5</b>	<b>More on nodal sets and partitions</b>	<b>37</b>
<b>6</b>	<b>Playing around Pleijel's argument.</b>	<b>40</b>
<b>7</b>	<b>Minimal partitions and Courant-sharp</b>	<b>43</b>
7.1	Preliminaries . . . . .	43
7.2	Definition of an exhausting family $N(u, \alpha)$ . . . . .	43
7.3	Proof of Theorem 1.17. . . . .	44
<b>8</b>	<b>Further properties of subpartitions</b>	<b>50</b>
<b>9</b>	<b>Example 1: The case of the disk</b>	<b>53</b>
<b>10</b>	<b>Example 2: the case of the rectangle</b>	<b>56</b>
<b>A</b>	<b>Zeros of Bessel functions</b>	<b>61</b>
<b>B</b>	<b>Alternative approach in the regular case</b>	<b>62</b>

# 1 Introduction and main results

We consider mainly two-dimensional Laplacians operators in bounded domains. We would like to analyze the relations between the nodal domains of the eigenfunctions of the Dirichlet Laplacians and the partitions by  $k$  open sets  $D_i$  which are optimal in the sense that the maximum over the  $D_i$ 's of the ground state energy of the Dirichlet realization of the Laplacian in  $D_i$  is minimal.

## 1.1 Definitions and notations

Let us consider a Schrödinger operator

$$H = -\Delta + V \tag{1.1}$$

on a bounded domain  $\Omega \subset \mathbb{R}^2$  with Dirichlet boundary condition.

In the whole article (except in Section 3), we will consider that  $\Omega$  satisfies the following condition of smoothness :

### Assumption 1.1

*$\Omega$  has compact and piecewise  $\mathcal{C}^{1,+}$  boundary, i.e. piecewise  $\mathcal{C}^{1,\alpha}$  for some  $\alpha > 0$ . Moreover  $\Omega$  satisfies the interior cone property.*

This allows a finite number of corners (and cracks) of opening  $\alpha\pi$  (defined in Section 2).

The other general assumption is that

### Assumption 1.2

*The potential  $V$  belongs to  $L^\infty(\Omega)$ .*

Under these assumptions (which will not be recalled at each statement),  $H$  is selfadjoint if viewed as the Friedrichs extension of the quadratic form associated to  $H$  with form domain  $W_0^{1,2}(\Omega)$  and form core  $\mathcal{C}_0^\infty(\Omega)$ . We denote  $H$  by  $H(\Omega)$ . We are interested in the eigenvalue problem for  $H(\Omega)$  and note that under our assumptions  $H(\Omega)$  has compact resolvent and its spectrum, which will be denoted by  $\sigma(H(\Omega))$  is discrete and consists of eigenvalues  $\{\lambda_k\}_{k=1}^\infty$  with finite multiplicities which tend to infinity, so that

$$\lambda_1 < \lambda_2 \leq \lambda_3 \leq \dots \leq \lambda_k \leq \dots \tag{1.2}$$

and such that the associated eigenfunctions  $u_k$  can be chosen to form an orthonormal basis for  $L^2(\Omega)$ .

Without loss of generality we can assume that the  $u_k$  are real valued and by elliptic regularity (see also Proposition 2.8 in Section 2) we have :

$$u_k \in \mathcal{C}^{1,\alpha}(\Omega) \cap \mathcal{C}_0^0(\overline{\Omega}) , \quad (1.3)$$

for any  $\alpha < 1$ .

We know that  $u_1$  can be chosen to be strictly positive in  $\Omega$ , but the other eigenfunctions  $u_k$  ( $k \geq 2$ ) must have zerosets. We define for any function  $u \in \mathcal{C}_0^0(\overline{\Omega})$

$$N(u) = \overline{\{x \in \Omega \mid u(x) = 0\}} \quad (1.4)$$

and call the components of  $\Omega \setminus N(u)$  the nodal domains of  $u$ . The number of nodal domains of such a function will be called  $\mu(u)$ .

We now introduce the notions of partition and minimal partition.

**Definition 1.3**

Let  $1 \leq k \in \mathbb{N}$ . We will call **partition** (or *k-partition* if we want to indicate the cardinality of the partition) of  $\Omega$  a family  $\mathcal{D} = \{D_i\}_{i=1}^k$  of mutually disjoint subsets of  $\Omega$  :

$$D_i \cap D_j = \emptyset , \forall i \neq j \text{ and } \cup_{i=1}^k D_i \subset \Omega . \quad (1.5)$$

We call it **open** if the  $D_i$  are open sets of  $\Omega$ , **connected** if the  $D_i$  are connected.

We denote by  $\mathfrak{D}_k$  the set of open connected *k-partitions* of  $\Omega$ .

We now introduce the notion of spectral minimal partition sequence.

**Definition 1.4**

Let  $H = H(\Omega)$  as above. For  $\mathcal{D}$  in  $\mathfrak{D}_k$ , we introduce

$$\Lambda(\mathcal{D}) = \max_i \lambda(D_i) , \quad (1.6)$$

where  $\lambda(D_i)$  is the ground state energy of  $H(D_j)$ .

**Remark 1.5**

When  $D$  is not sufficiently regular, we define  $\lambda(D)$  differently. See Definition 3.1.

**Definition 1.6**

For any integer  $k \geq 1$ , we define

$$\mathfrak{L}_k = \inf_{\mathcal{D} \in \mathfrak{D}_k} \Lambda(\mathcal{D}). \quad (1.7)$$

We call the sequence  $\{\mathfrak{L}_k\}_{k \geq 1}$  the spectral minimal partition sequence of  $H(\Omega)$ .

For given  $k$ , we call  $k$ -minimal partition a partition  $\mathcal{D} \in \mathfrak{D}_k$ , such that  $\mathfrak{L}_k = \Lambda(\mathcal{D})$ .

**Remark 1.7**

If  $k = 2$ , it is rather well known (see for example [16] or [9]) that  $\mathfrak{L}_2$  is the second eigenvalue and the associated minimal 2-partition is the nodal partition associated to the second eigenfunction.

We now introduce the notion of strong partition.

**Definition 1.8**

A partition  $\mathcal{D} = \{D_i\}_{i=1}^k$  of  $\Omega$  in  $\mathfrak{D}_k$  is called **strong** if

$$\text{Int}(\overline{\cup_i D_i}) \setminus \partial\Omega = \Omega. \quad (1.8)$$

Attached to a partition, we can naturally associate a closed set in  $\overline{\Omega}$  defined by

$$N(\mathcal{D}) = \overline{\cup_i (\partial D_i \cap \Omega)}. \quad (1.9)$$

This leads us to introduce the definition of a regular closed set. This definition is modeled on some (but not all) of the properties of the nodal set of an eigenfunction of a Schrödinger operator (see Section 2).

**Definition 1.9**

A closed set  $N \subset \overline{\Omega}$  is regular (and we write in this case belongs to  $N \in \mathcal{M}(\Omega)$  (and we say in this case that it is regular) if  $N$  meets the following requirements:

(i)

There are finitely many distinct  $x_i \in \Omega \cap N$  and associated positive integers  $\nu_i$  with  $\nu_i \geq 2$  such that, in a sufficiently small neighborhood of each of the  $x_i$ ,  $N$  is the union of  $\nu_i(x_i)$   $\mathcal{C}^{1,+}$  curves (non self-crossing) with one end at  $x_i$  (and each pair defining at  $x_i$  a positive angle in  $(0, 2\pi)$ ) and such that in the complement of these points in  $\Omega$ ,  $N$  is locally diffeomorphic to a  $\mathcal{C}^{1,1-}$  (i.e.  $\mathcal{C}^{1,\alpha}$  for any  $\alpha \in ]0, 1[$ ) curve.

(ii)

$\partial\Omega \cap N$  consists of a (possibly empty) finite set of points  $z_i$ , such that, at each  $z_i$ ,  $\rho_i \mathcal{C}^{1,+}$  half-lines belonging to  $N$  (with  $\rho_i \geq 1$ ) hit the boundary.

(iii)

Moreover the half curves meet with equal angle at each critical point of  $N \cap \Omega$  and also at each point of  $N \cap \partial\Omega$  together with the boundary.

Complementarily, we introduce the notion of regular partition.

**Definition 1.10**

A strong partition  $\mathcal{D}$  is regular (and we write in this case  $\mathcal{D} \in \mathcal{R}(\Omega)$ ) if there exists a regular closed set  $N$  such that  $\mathcal{D} = \mathcal{D}(N)$ , where  $\mathcal{D}(N)$  is the family of the connected components of  $\Omega \setminus N$  belongs (by definition) to  $\mathcal{R}(\Omega)$ .

## 1.2 Main results

Although some of the statements could be obtained under weaker assumptions we assume below that  $\Omega$  is bounded and connected.

It has been proved<sup>1</sup> by Conti-Terracini-Verzini [7, 8, 9] that

**Theorem 1.11**

For any  $k$ , there exists a  $k$ -minimal regular strong partition.

The first aim of this paper is to show the

**Theorem 1.12**

Any  $k$ -minimal partition has a connected, regular and strong representative.

Here we need to explain what we mean by representative (which involves implicitly the notion of capacity). This involves indeed a notion of equivalence classes. Two  $k$ -partitions  $\mathcal{D}$  and  $\tilde{\mathcal{D}}$  are equivalent if there is a labeling such that for any ground state  $u_i$  associated of  $D_i$ , there is a ground state  $\tilde{u}_i$  associated to  $\tilde{D}_i$  such that  $u_i = \tilde{u}_i$  in  $W_0^1(\Omega)$ , and conversely.

Once this notion is introduced it is natural to look for the existence of a regular representative and uniqueness will always be inside this class.

---

<sup>1</sup>But these papers treat only smoother boundaries than assumed in the whole article. So we will prove here a slight generalization.

In general, there is no reason for a minimal partition to be unique (and here we speak of uniqueness of equivalence classes). This can for example occur in presence of symmetries. However, we will show that the uniqueness property always holds for subpartitions of a given minimal partition. More precisely, we have

**Theorem 1.13**

Let  $\mathcal{D}$  be a minimal  $k$ -partition relative to  $\mathfrak{L}_k(\Omega)$ . Let  $\mathcal{D}' \subset \mathcal{D}$  be a subpartition of  $\mathcal{D}$  into  $1 \leq k' < k$  elements and assume that

$$\Omega' := \text{Int} \left( \bigcup_{D_i \in \mathcal{D}'} \overline{D_i} \right),$$

is connected. Then  $\mathfrak{L}_k(\Omega) = \mathfrak{L}_{k'}(\Omega')$  and the  $k'$ -minimal partition of  $\Omega'$  is unique.

A natural question is whether a minimal partition is the nodal partition induced by an eigenfunction. Theorem 1.14 gives a simple criterion for a partition to be associated to a nodal set. For this we need some additional definitions.

We say that  $D_i, D_j$  are **neighbors** and we write  $D_i \sim D_j$ , if the set  $D_{i,j} := \text{Int}(\overline{D_i \cup D_j}) \setminus \partial\Omega$  is connected. We then construct for each  $\mathcal{D}$  a graph  $G(\mathcal{D})$  by associating to each  $D_i$  a vertex and to each pair of neighbors  $(D_i, D_j)$  an edge. This is an undirected graph without multiple edges or loops. We will say that the graph is **bipartite** if it can be colored by two colors. We recall that the graph associated to a collection of nodal domains of an eigenfunction is always bipartite. In this case, we say that the partition is **admissible**. We have now the following converse theorem :

**Theorem 1.14**

If the graph of the minimal partition of  $\Omega$  is bipartite, this is a partition associated to the nodal set of an eigenfunction of  $H(\Omega)$  corresponding to  $\mathfrak{L}_k(\Omega)$ .

This theorem was already obtained in [16] by adding a strong a priori regularity assumption on the partition and the assumption that  $\Omega$  is simply connected. Any subpartition of cardinality two  $(D_i, D_j)$  corresponds indeed to a nodal partition of some eigenfunction associated to the second eigenvalue of  $H(D_{i,j})$ . This implies the Pair Compatibility Condition (see in Appendix B) and Theorem B.1 can be applied.

The proof given here is more general (but more difficult) and is actually a

byproduct of the proof of Theorem 1.12, which will directly give an eigenfunction whose nodal domains form the partition.

A natural question is now to determine how general is the situation described in the previous theorem. The surprise is that this will only occur in the so called Courant-sharp situation. Before stating precisely our second main result we need to introduce some further statements and notations. The Courant Nodal Theorem says :

**Theorem 1.15**

*Let  $k \geq 1$ ,  $\lambda_k = \lambda_k(\Omega)$  the  $k$ -th eigenvalue of  $H(\Omega)$  and  $u$  any real associated eigenfunction. Then the number of nodal domains  $\mu(u)$  of  $u$  satisfies  $\mu(u) \leq k$ .*

When the number of nodal domains  $\mu(u)$  satisfies

$$\mu(u) = k ,$$

we will say, as in [4], that  $u$  is **Courant-sharp**.

**Definition 1.16**

*For any integer  $k \geq 1$ , we denote by  $L_k(\Omega)$  (or simply  $L_k$ ) the smallest eigenvalue whose eigenspace contains an eigenfunction with  $k$  nodal domains.*

In general, we will show in Corollary 5.6, that

$$\lambda_k(\Omega) \leq \mathfrak{L}_k(\Omega) \leq L_k(\Omega) . \tag{1.10}$$

The last goal consists in giving the full picture of the equality cases :

**Theorem 1.17**

*If  $\mathfrak{L}_k(\Omega) = L_k(\Omega)$  or  $\mathfrak{L}_k(\Omega) = \lambda_k(\Omega)$ , then*

$$\lambda_k(\Omega) = \mathfrak{L}_k(\Omega) = L_k(\Omega) .$$

*In addition, one can find in the eigenspace associated to  $\lambda_k$  an eigenfunction  $u$  such that  $\mu(u) = k$ .*

In other words, the only case when the  $k$ - nodal domains of an eigenfunction of  $H(\Omega)$  form a minimal partition is the case when this eigenfunction is Courant-sharp.

### 1.3 Organization of the paper

The paper is organized as follows. We first start in Section 2 by recalling and extending (at the boundary) results on the local properties of the nodal set of an eigenfunction. Section 3 is devoted to the analysis of the geometrical properties of minimal partitions in  $\mathbb{R}^N$ . Section 4 gives stronger results but limited to the two-dimensional case, which is our main subject. This gives in particular the proof of our first Main Theorem 1.12. Sections 5 and 6 are devoted to additional properties of the minimal partitions. We discuss different notions related to the spectrum and revisit Pleijel's theorem and its proof. Section 7 gives the proof of the second Main Theorem 1.17 permitting to show that when a  $k$ -minimal partition is a nodal family then the corresponding eigenvalue is the  $k$ -th one. In Section 8, we complete the proofs and the statements concerning subpartitions. In Sections 9 and 10 we analyze in great detail the various spectra of specific  $H(\Omega)$  in connection with minimal partitions. This leads in particular to nice conjectures and open problems. Finally, we develop in two appendices useful results which will complete some proofs or help the reader.

#### Acknowledgements

The two first authors are supported by the European Research Network 'Post-doctoral Training Program in Mathematical Analysis of Large Quantum Systems' with contract number HPRN-CT-2002-00277, and the ESF Scientific Programme in Spectral Theory and Partial Differential Equations (SPECT). The support of the ESI was also very useful.

The two first authors would also like to thank M. Van den Berg who was stimulating their interest for this question, M. Dauge for discussions on problems with corners and V. Bonnaillie-Noel and G. Vial for their help in the analysis of the problem via numerical computations. The third author was partially supported by Italian MIUR, national project "Variational Methods and Non-linear Differential Equations". She wishes to thank ESI, where a part of this work was done, for the support and the kind hospitality.

## 2 Preliminaries: Hölder regularity of nodal sets

It is a well known property of nodal sets of eigenfunctions to be the union of curves ending either at interior singular points or at the boundary. This

section is devoted to the analysis of the regularity of the nodal curves in the Hölder spaces  $\mathcal{C}^{1,\varepsilon}$ . A word of caution must be entered at this point: with regularity we mean *global regularity* of the nodal branch up to the singularities or the boundary. This is not a completely obvious issue (basically because of the lack of regularity of our solutions and, possibly, of the boundary of the domain) and will require a revisitation of the well known asymptotic estimates about critical points of eigenfunctions. To start with, we recall the classical local regularity result by Hartman and Wintner ([15], Corollary 1), stating that interior critical points of non zero solutions to our class of equations are isolated and have finite (local) multiplicity  $m$ . In addition the solution satisfies, for some  $c \neq 0$ , the asymptotic formula

$$u(r, \theta) = cr^{m+1} \cos((m+1)(\theta + \theta_0)) + o(r^{m+1}), \quad r = |z - z_0|. \quad (2.1)$$

Here we identify  $\mathbb{R}^2$  to  $\mathbb{C}$  and use either  $z$ , or  $(x, y)$ , or  $(r, \theta)$  for a point of  $\mathbb{R}^2$ , with the standard notations :

$$z = r \exp i\theta, \quad z = x + iy, \quad x = r \cos \theta, \quad y = r \sin \theta.$$

We shall need a refined version of it which is stated below:

**Theorem 2.1**

Let  $\Omega$  be open and  $V \in L^\infty(\Omega)$ . Assume  $u \in W_{\text{loc}}^{1,2}(\Omega)$  solves

$$-\Delta u + V(x, y)u = 0,$$

in the distributional sense.

Let  $z_0 = (x_0, y_0) \in \Omega$  be such that  $u(x_0, y_0) = 0$  and  $\nabla u(x_0, y_0) = 0$ ; then, in a neighbourhood of  $z_0$ ,

- (a) There are an integer  $n$ , a complex-valued function  $\xi$  of class  $\mathcal{C}^{0,+}$  such that  $\xi(z_0) \neq 0$  and

$$u_x + iu_y = r^n e^{-in\theta} \xi(x, y), \quad r = |z - z_0| \quad (2.2)$$

- (b) There is a function  $\tilde{\xi}$  of class  $\mathcal{C}^{0,+}$  such that  $\tilde{\xi}(x_0, y_0) = 0$  and

$$u(x, y) = \frac{r^{n+1}}{n+1} \left( \Re(\xi(x_0, y_0)) \cos(n+1)\theta + \Im(\xi(x_0, y_0)) \sin(n+1)\theta + \tilde{\xi}(x, y) \right). \quad (2.3)$$

- (c) There exists a positive radius  $R$  such that  $u^{-1}(\{0\}) \cap B(z_0, R)$  is composed by  $2n$   $\mathcal{C}^{1,+}$ -simple arcs which all end in  $z_0$  and whose tangent lines at  $z_0$  divide the disc into  $2n$  angles of equal amplitude.

**Proof :**

Following the paper by Hartman and Wintner [15], denote  $w = u_y + iu_x$  and set  $z_0 = 0$ . Then, it is shown there that, if,

$$u = o(|z|^k), \quad (2.4)$$

for some integer  $k \geq 0$ , then the Cauchy formula is available:

$$2\pi i \frac{w(\zeta)}{\zeta^k} = \int_{|z|=R} \frac{w(z)}{z^k(z-\zeta)} dz - \int_{|z|<R} \frac{V(z)u(z)}{z^k(z-\zeta)} dx dy. \quad (2.5)$$

where  $R > 0$  is fixed and the double integral over the disk is absolutely convergent. We now wish to show that it follows that the left-hand side is Hölder continuous in  $\zeta$  in a neighbourhood of the origin. The line integral is smooth in  $\zeta$ , since the integrand has no singularities on the circle. Concerning the second term, notice that we can find a constant  $K$  such that

$$\begin{aligned} \left| \int_{|z|<R} \frac{V(z)u(z)}{z^k} \left( \frac{1}{z-\zeta_1} - \frac{1}{z-\zeta_2} \right) dx dy \right| \leq \\ \int_{|z|<R} \frac{|V(z)u(z)|}{|z|^k} \left| \frac{|\zeta_1 - \zeta_2|}{|z-\zeta_1||z-\zeta_2|} \right| dx dy \leq \\ K|\zeta_1 - \zeta_2| |\log |\zeta_1 - \zeta_2||. \end{aligned}$$

Now we wish to show that (2.4) can not be verified for every integer. To this aim, we integrate Equation (2.5) over the disk and, taking absolute values, we obtain:

$$2\pi \int_{|z|<R} \frac{|w(z)|}{|z|^k} dx dy \leq 2\pi R \int_{|z|=R} \frac{|w(z)|}{|z|^k} |dz| + 2\pi R \int_{|z|<R} \frac{|V(z)||u(z)|}{|z|^k} dx dy. \quad (2.6)$$

Following [15], and using the identity

$$u(r, \theta) = \int_0^r (u_x(\rho, \theta) \cos \theta + u_y(\rho, \theta) \sin \theta) d\rho, \quad (2.7)$$

we observe that :

$$|u(z)| \leq \int_0^1 |zw(tz)| dt,$$

implies

$$\int_{|z|<R} \frac{|V(z)||u(z)|}{|z|^k} dx dy \leq K \int_{|z|<R} \frac{|zw(z)|}{|z|^k} dx dy \leq KR \int_{|z|<R} \frac{|w(z)|}{|z|^k} dx dy.$$

Thus, for  $R$  sufficiently small, inequality (2.6) leads to

$$\int_{|z|<R} \frac{|w(z)|}{|z|^k} dx dy \leq 2R \int_{|z|=R} \frac{|w(z)|}{|z|^k} |dz|. \quad (2.8)$$

We have now fixed  $R > 0$  such that (2.8) is satisfied. Let us assume that  $w(z_0) \neq 0$  for some  $|z_0| < R$ . Then, for a constant independent of  $k$  there holds

$$|w(z_0)| \leq K \left( \frac{|z_0|}{R} \right)^k, \quad k = 1, 2, \dots$$

Let us take the limit  $k \rightarrow +\infty$  in this inequality. Then the limit of the sequence  $(|z_0|/R)^k$  does not vanish, in contradiction with  $|z_0| < R$ . This completes the proof of point (a) in the statement of the Theorem. Point (b) follows from point (a) together with the identity (2.7).

To prove point (c) we choose a branch of the nodal set and we choose, as a regular parametrization the path  $z(t) = r(t)e^{i\theta(t)}$ , where the pair  $(r(t), \theta(t))$  solves the following system of ordinary differential equations:

$$\begin{cases} \dot{r} = \frac{1}{r^{n+1}} (xu_y - yu_x) \\ \dot{\theta} = \frac{1}{r^{n+2}} (xu_x + yu_y) \end{cases},$$

where  $\dot{r}$  and  $\dot{\theta}$  denote respectively the derivative of  $r(t)$  and  $\theta(t)$  with respect to  $t$ .

One can easily prove by the points (a) and (b) that both functions  $t \mapsto \dot{r}(t)$  and  $t \mapsto r(t)\dot{\theta}(t)$  are Hölder continuous; therefore both  $r$  and  $\theta$  are Hölder continuous functions. Hence they can be extended through the singularity. Since the parametrization is regular ( $\dot{z} \neq 0$ ), the assertion follows from the equation

$$\dot{z} = \dot{r}(t)e^{i\theta(t)} + ir(t)\dot{\theta}(t)e^{i\theta(t)}.$$

■

### Remark 2.2

*Theorem 2.1 extends, with the same argument, to the case when the potential  $V$  has a singularity at  $z_0$ , provided there exists  $\beta < 1$  and  $K$  such that*

$$|V(x, y)| \leq \frac{K}{|z - z_0|^\beta}.$$

*This fact will be useful when we shall consider the case of domains with corners or cracks; indeed such singular potentials result as conformal factors*

associated with the complex exponentials.

Note that this singular situation was also analyzed, but for the interior problem, in [19] and [20] and that in this case the authors obtain a better regularity.

In order to examine the regularity up to the boundary of the nodal partition associated to an eigenfunction we now extend a known result by Alessandrini [3] (which treats the convex case) to our setting. The proof exploits the classical Kellog–Warschawski theorem on the boundary regularity of conformal mappings which states that any conformal map on a  $\mathcal{C}^{1,\varepsilon}$  domain extends continuously on the boundary keeping the same regularity (see the book by Pommerenke [24], Theorem 3.6 in Chapter 3).

**Theorem 2.3**

Let  $\varepsilon > 0$  and  $\Omega$  be an open set with  $\mathcal{C}^{1,\varepsilon}$  boundary and  $V \in L^\infty(\Omega)$ . Assume  $u \in W_0^{1,2}(\Omega)$  solves

$$-\Delta u + V(x, y)u = 0 ,$$

in the distributional sense. Then the associated nodal partition is regular. More precisely if  $\overline{u^{-1}(\{0\})}$  intersects  $\partial\Omega$  at  $z_0$ , then there exist an integer  $m$  and  $R > 0$  such that  $u^{-1}(\{0\}) \cap B(z_0, R)$  is composed by  $m$   $\mathcal{C}^{1,\varepsilon}$ -simple arcs which all end in  $z_0$  and whose tangent lines at  $z_0$  divide the tangent cone  $\Gamma(z_0)$  into  $m + 1$  angles of equal opening.

**Proof :**

The result immediately follows from Theorem 2.1 in the case of the half-plane: indeed one can extend  $u$  by a reflection to the other half-plane and reduce to the case of the interior zeros. The general case reduces to that of the half-space through the Riemann mapping Theorem. Indeed, by [24] (Theorem 3.6 in Chapter 3) the Hölder regularity  $\mathcal{C}^{1,\varepsilon}$  of  $\partial\Omega$  implies the same regularity property for the extensions, up to boundary, of the Riemann map  $f$  and of its inverse. Since the composition of  $\mathcal{C}^{1,\varepsilon}$  maps enjoys the same regularity property, the statement follows. ■

Now we wish to extend Theorem 2.3 to the case of domains possessing corners or cracks. To be precise we start with the following

**Definition 2.4**

Let  $\varepsilon \in ]0, 1]$ . We say that  $\partial\Omega$  has a  $\mathcal{C}^{1,\varepsilon}$ -corner of opening  $\alpha\pi$  ( $0 \leq \alpha \leq 2$ ) at  $z_0$  if, in a sufficiently small neighborhood,  $\partial\Omega$  contains the union of two curves of class  $\mathcal{C}^{1,\varepsilon}$  (non self-crossing) ending at  $z_0$ , and such that  $\Omega$  lies in

the curvilinear sector of angle opening  $\alpha\pi$  spanned by the two arcs, which does not intersect other components of the boundary  $\partial\Omega$ .

**Remark 2.5**

Note that the boundary  $\partial\Omega$  can have several corners of angle opening  $\alpha_i$  at the same point  $z_0$ : of course the sum of all the angles does not exceed  $2\pi$ . Moreover, it is worthwhile noticing that we allow the presence of cracks (i.e. corners of angle opening  $2\pi$  where the two curves coincide), exterior cusps (i.e. corners of angle opening  $2\pi$  spanned by two distinct curves), as well as angles of any possible positive angles, (positiveness is required by the interior cone property). Finally, a corner can be a point of smoothness of the boundary, when its angle opening is  $\pi$ .

Our next goal is to prove the following result

**Theorem 2.6**

Let  $V \in L^\infty(\Omega)$  and  $\varepsilon \in ]0, 1]$ . Assume  $u \in W_0^{1,2}(\Omega)$  solves  $-\Delta u + V(x, y)u = 0$  in the distributional sense, in a neighbourhood of some  $z_0 \in \partial\Omega$ , a  $\mathcal{C}^{1,\varepsilon}$ -corner of opening  $\alpha\pi$  ( $0 < \alpha \leq 2$ ). If  $\overline{u^{-1}(\{0\})}$  intersects  $\partial\Omega$  at  $z_0$ , then there exist an integer  $m$  and  $R > 0$  such that  $u^{-1}(\{0\}) \cap B(z_0, R)$  is composed by  $m$   $\mathcal{C}^{1,\varepsilon'(\alpha)}$ -simple arcs which all end at  $z_0$  and whose tangent lines at  $z_0$  divide the tangent cone  $\Gamma(z_0)$  into  $m + 1$  angles of equal amplitude. In addition

$$\varepsilon'(\alpha) = \begin{cases} \varepsilon \min(\alpha, 1/\alpha) & \text{if } 1/2 < \alpha \leq 2 \\ 2^n \varepsilon \alpha & \text{if } 1/2^{(n+1)} < \alpha \leq 1/2^n \end{cases}$$

To prove the theorem we shall first straighten the corner and then apply Theorem 2.3. We shall need the following basic result.

**Proposition 2.7**

Let  $\varepsilon \in ]0, 1]$  and let  $C$  be a Hölder-continuous arc ending at the origin, without self-intersections. Let  $w(\tau)$ ,  $\tau \in [0, \bar{\tau}]$  be a regular parametrization of  $C$  such that  $w(0) = 0$  and  $w'(0) \neq 0$ , and define the curve  $\mathcal{C}^{1/\alpha}$  by the parametrization

$$t \mapsto v(t) := (w(t^\alpha))^{1/\alpha}.$$

Then, for any  $\alpha > 0$

$$C \in \mathcal{C}^{1,\varepsilon} \implies \mathcal{C}^{1/\alpha} \in \mathcal{C}^{1,\varepsilon \min(1,\alpha)}$$

**Proof :**

We have

$$v'(t) = w'(t^\alpha)u(t^\alpha), \quad u(\tau) := (\tau w(\tau))^{-1+1/\alpha}.$$

Obviously  $v$  defines a regular parametrization of  $C^{1/\alpha}$ . At first we remark that  $u$  is Hölder continuous with exponent  $\varepsilon$ . Indeed

$$\left| \frac{w(\tau_1)}{\tau_1} - \frac{w(\tau_2)}{\tau_2} \right| = \left| \int_0^1 (w'(\tau_2 s) - w'(\tau_1 s)) ds \right| \leq K |\tau_1 - \tau_2|^\varepsilon.$$

Therefore the product  $w'(\tau)u(\tau)$  is of the same class  $\mathcal{C}^{0,\varepsilon}$  and the composition  $v'(t) = w'(t^\alpha)u(t^\alpha)$  is in the Hölder space  $\mathcal{C}^{0,\varepsilon \min(1,\alpha)}$ . ■

**Proof :**

We are in position to prove Theorem 2.6. We consider separately the two cases :

First, we assume the case of openings satisfying the inequality  $1/2 < \alpha \leq 2$ . We straighten the corner as in Proposition 2.7, by the map  $z \rightarrow z^{1/\alpha}$ . Then, through this composition, the boundary  $\partial\Omega^{1/\alpha}$  becomes smooth (of class  $\mathcal{C}^{1,\varepsilon \min(1,\alpha)}$ ) while the potential  $V$  has to be multiplied by the conformal factor  $2\alpha^2|z - z_0|^{2(\alpha-1)}$ , which is singular whenever  $\alpha < 1$ . As already observed in Remark 2.2, this is not a problem if  $\alpha > 1/2$ . Thanks to Theorem 2.3, the nodal set of the composition  $z \mapsto u(z^\alpha)$  is the union of arcs of class  $\mathcal{C}^{1,\varepsilon \min(1,\alpha)}$ . Now we take its inverse image through the map  $z \rightarrow z^\alpha$  and, applying again Proposition 2.7 we obtain the desired value of  $\varepsilon'(\alpha)$ .

Next we turn to the case when the opening is too small, that is when  $1/2^{(n+1)} < \alpha \leq 1/2^n$ , for some  $n \geq 1$ . Using again Theorem 3.6 in Chapter 3 of [24], one can easily construct, locally in  $\Omega$ , a conformal map of class  $\mathcal{C}^{1,\varepsilon}$  up to the boundary such that the image of one of the two arcs is a straight segment. Next step is to reflect the domain about this line and extend the function on the reflected corner, in such a way to double the opening, which is now  $2\alpha$ . In this procedure, the second arc, being composed with a  $\mathcal{C}^{1,\varepsilon}$  map, still remains in the same Hölder class. We iterate this reflection procedure  $n$  times, until  $1/2 < 2^n\alpha \leq 1$ , and we afterward proceed as in the proof of the case  $\alpha > 1/2$ . ■

Using the same technique of straightening the angles by conformal maps, one can easily prove the following

**Proposition 2.8**

*Let  $V \in L^\infty(\Omega)$  and  $\varepsilon \in ]0, 1]$ . Assume  $u \in W_0^{1,2}$  solves  $-\Delta u + V(x, y)u = 0$  in the distributional sense, in a neighbourhood of some  $z_0 \in \partial\Omega$ , a  $\mathcal{C}^{1,\varepsilon}$ -corner of opening  $\alpha\pi$  ( $0 < \alpha \leq 2$ ). Then, if  $\alpha \leq 1$ ,  $u \in \mathcal{C}^{1,\varepsilon}$ , locally at  $z_0$ ; otherwise, if  $1 < \alpha \leq 2$ , we only have  $u \in \mathcal{C}^{0,1/\alpha}$ .*

### 3 Optimal partitions in $N$ dimensions

Let  $\Omega \subset \mathbb{R}^N$  be a connected, open bounded domain. Note that at this stage we don't need any regularity of the boundary. However we will need it later (in Section 4), in the 2-dimension case in order to describe the local structure of nodal lines at their intersection with the boundary.

#### Definition 3.1

For any measurable  $D \subset \Omega$  and for  $V \in L^\infty(\Omega)$ , let  $\lambda_1(D)$  denotes the first eigenvalue of the Dirichlet realization of the Schrödinger operator in the following generalized sense. We define

$$\lambda_1(D) = +\infty ,$$

if  $\{u \in W_0^1(\Omega), u \equiv 0 \text{ a.e. on } \Omega \setminus D\} = \{0\}$ , and

$$\lambda_1(D) = \min \left\{ \frac{\int_{\Omega} (|\nabla u(x)|^2 + V(x)u(x)^2) dx}{\int_{\Omega} |u(x)|^2 dx} : u \in W_0^1(\Omega) \setminus \{0\}, u \equiv 0 \text{ a.e. on } \Omega \setminus D \right\} ,$$

otherwise. We call groundstate any function  $\phi$  achieving the above minimum.

We shall always assume that

$$\lambda_1(\Omega) > 0 .$$

#### Remark 3.2

The presence of an  $L^\infty$  potential  $V$  does not create particular problems. We prefer to simplify the notations to explain all the proofs with the additional assumption that  $V$  is identically 0. In this case the positivity on  $\lambda_1(\Omega)$  is effectively satisfied. In the general case, we can always assume this property by adding a constant to  $V$ .

We observe that the minimization problem always possesses a (possibly not unique) non negative solution  $\phi \geq 0$ . We shall always make this choice. Next we consider the following class of minimal partition problems :

$$\mathfrak{L}_{k,p} := \inf_{\mathcal{B}_k} \left( \frac{1}{k} \sum_{i=1}^k (\lambda_1(D_i))^p \right)^{1/p} , \quad (3.1)$$

$$\mathfrak{L}_k := \inf_{\mathcal{B}_k} \max_{i=1,\dots,k} (\lambda_1(D_i)) \quad (3.2)$$

where the minimization is taken over the class of partitions in  $k$  “disjoint” measurable subsets of  $\Omega$

$$\mathcal{B}_k := \left\{ \mathcal{D} = (D_1, \dots, D_k) : \bigcup_{i=1}^k D_i \subset \Omega, |D_i \cap D_j| = 0 \text{ if } i \neq j \right\}.$$

**Remark 3.3**

The values  $\mathfrak{L}_k$  considered in this section can be viewed as a relaxation of those defined in the introduction. We have indeed replaced “open” by “measurable”. We keep the same notation, for we shall prove as a part of our regularity theory that, in all the interesting cases, the two definitions coincide.

The main result of this section is the following

**Theorem 3.4**

Let  $\mathcal{D} = (\tilde{D}_1, \dots, \tilde{D}_k) \in \mathcal{B}_k$  be any minimal partition associated with  $\mathfrak{L}_k$  and let  $(\tilde{\phi}_i)_i$  be any set of positive eigenfunctions normalized in  $L^2$  corresponding to  $(\lambda_1(\tilde{D}_i))_i$ . Then, there exist  $a_i \geq 0$ , not all vanishing, such that the functions  $\tilde{u}_i = a_i \tilde{\phi}_i$  verify in  $\Omega$  the differential inequalities in the distributional sense

$$(I1) \quad -\Delta \tilde{u}_i \leq \mathfrak{L}_k \tilde{u}_i, \quad \forall i = 1, \dots, k,$$

$$(I2) \quad -\Delta \left( \tilde{u}_i - \sum_{j \neq i} \tilde{u}_j \right) \geq \mathfrak{L}_k \left( \tilde{u}_i - \sum_{j \neq i} \tilde{u}_j \right), \quad \forall i = 1, \dots, k.$$

**Remark 3.5**

Note that at this stage we do not know if the  $\tilde{D}_i$ 's are connected and consequently if the  $\tilde{\phi}_i$ 's are unique. It will be shown in the next section that these properties are true in two dimensions.

The following results were proved in [9]:

**Theorem 3.6**

Let  $p \in [1, +\infty)$  and let  $\mathcal{D} = (D_1, \dots, D_k) \in \mathcal{B}_k$  be a minimal partition associated with  $\mathfrak{L}_{k,p}$  and let  $(\phi_i)_i$  be any set of positive eigenfunctions normalized in  $L^2$  corresponding to  $(\lambda_1(D_i))_i$ . Then, there exist  $a_i > 0$ , such that the functions  $u_i = a_i \phi_i$  satisfy in  $\Omega$  the differential inequalities in the distribution sense

$$(I1) \quad -\Delta u_i \leq \lambda_1(D_i) u_i,$$

$$(I2) \quad -\Delta \left( u_i - \sum_{j \neq i} u_j \right) \geq \lambda_1(D_i) u_i - \sum_{j \neq i} \lambda_1(D_j) u_j.$$

**Remark 3.7**

In particular, this implies that  $U = (u_1, \dots, u_k)$  is in the class  $\mathcal{S}^*$  as defined in [8]. Hence Theorem 8.3 in [8] ensures the Lipschitz continuity of the  $u_i$ 's in the interior of  $\Omega$ . Therefore we can choose a partition made of open representatives  $D_i = \{u_i > 0\}$ .

Moreover, taking the limit as  $p \rightarrow +\infty$ , the following result was shown in [9]:

**Theorem 3.8**

There holds

$$\lim_{p \rightarrow +\infty} \mathfrak{L}_{k,p} = \mathfrak{L}_k .$$

Moreover, there exists a minimizer of  $\mathfrak{L}_k$  such that (I1)- (I2) hold for suitable non negative multiples  $u_i = a_i \phi_i$  of an appropriate set of associated eigenfunctions.

**Let us start the proof of Theorem 3.4.**

Let  $(\tilde{D}_1, \dots, \tilde{D}_k) \in \mathcal{B}_k$  be a particular minimal partition associated with  $\mathfrak{L}_k$  and let  $(\tilde{\phi}_1, \dots, \tilde{\phi}_k)$  be any choice of associated eigenfunctions. We wish to prove that (I1)-(I2) hold for a suitable set of multiples of the  $\tilde{\phi}_j$ 's. We consider, for a given

$$q \in (1, N/(N-2)) , \quad (3.3)$$

(or  $q \in (1, +\infty)$  when  $N = 2$ ), the penalized Rayleigh quotient:

$$\mathcal{F}_{k,p}(u_1, \dots, u_k) = \left( \frac{1}{k} \sum_{i=1}^k \left( \frac{\int_{\Omega} |\nabla u_i(x)|^2 dx}{\int_{\Omega} |u_i(x)|^2 dx} \right)^p \right)^{1/p} + \sum_{i=1}^k \left( 1 - \frac{\int_{\Omega} u_i(x)^q \tilde{\phi}_i(x)^q dx}{\left( \int_{\Omega} u_i(x)^{2q} dx \int_{\Omega} \tilde{\phi}_i(x)^{2q} dx \right)^{1/2}} \right) .$$

We consider the minimization problem

$$\mathcal{M}_{k,p} = \inf \{ \mathcal{F}_{k,p}(u_1, \dots, u_k) : (u_1, \dots, u_k) \in \mathcal{U} \} , \quad (3.4)$$

where

$$\mathcal{U} = \{ (u_1, \dots, u_k) \in (W_0^1(\Omega))^k : u_i \cdot u_j = 0 , \text{ for } i \neq j , u_i \geq 0 , u_i \not\equiv 0 , \forall i = 1, \dots, k \} . \quad (3.5)$$

We note that the condition on  $q$  permits to have (weak and strong) continuity and differentiability in  $W_0^1(\Omega)$  of the penalization term, which involves integrals of powers of  $u_i$ . This will be used later to apply the direct method

of the Calculus of Variations and to differentiate  $\mathcal{F}_{k,p}$  at the minimum. It is also worthwhile noticing that  $\mathcal{F}_{k,p}$  is invariant by multiplication:

$$\mathcal{F}_{k,p}(a_1 u_1, \dots, a_k u_k) = \mathcal{F}_{k,p}(u_1, \dots, u_k), \forall a_i \neq 0. \quad (3.6)$$

Recalling Definition 3.1 we have:

**Proposition 3.9**

There holds, for every  $p \in [1, +\infty)$ ,

$$\frac{1}{k^{1/p}} \mathfrak{L}_k \leq \mathfrak{L}_{k,p} \leq \mathcal{M}_{k,p} \leq \mathfrak{L}_k.$$

**Proof**

It is an immediate consequence of Jensen and Hölder inequalities.

**Lemma 3.10**

For every  $p \in [1, +\infty)$ , the value  $\mathcal{M}_{k,p}$  is achieved.

**Proof**

Using the invariance by multiplication (3.6), we can choose a bounded minimizing sequence, having as weak limit the configuration  $(u_1, \dots, u_k) \in \mathcal{U}$ . Now the assertion simply follows from the weak lower semi-continuity of the norm and the compact embeddings of  $W_0^1(\Omega)$  into  $L^s(\Omega)$  for any  $s \in [1, +\infty)$ , whenever  $N = 2$ , and for any  $s \in [1, 2N/(N - 2))$  when  $N \geq 3$ .

**Lemma 3.11**

Let  $\Lambda > 0$  and let  $U = (u_1, \dots, u_k)$  be any minimizer of  $\mathcal{M}_{k,p}$  normalized in such a way that

$$\left( \int_{\Omega} |\nabla u_i|^2 dx \right)^{p-1} = \left( \Lambda \int_{\Omega} |u_i|^2 dx \right)^p, \quad \forall i = 1, \dots, k. \quad (3.7)$$

Define

$$f_i(u)(x) = \frac{-\gamma q}{2 \left( \int_{\Omega} u(x)^{2q} dx \int_{\Omega} \tilde{\phi}_i(x)^{2q} dx \right)^{1/2}} \left[ u(x)^{q-1} \tilde{\phi}_i(x)^q - \frac{\int_{\Omega} u(x)^q \tilde{\phi}_i(x)^q dx}{\int_{\Omega} u(x)^{2q} dx} u(x)^{2q-1} \right], \quad (3.8)$$

where

$$\gamma = \Lambda^{-p} \left( \frac{1}{k} \sum_{i=1}^k \left( \frac{\int_{\Omega} |\nabla u_i(x)|^2 dx}{\int_{\Omega} |u_i(x)|^2 dx} \right)^p \right)^{1-1/p}. \quad (3.9)$$

Then  $U$  satisfies the differential inequalities in the distribution sense

$$(I1) \quad -\Delta u_i \leq \lambda_1(D_i)u_i + f_i(u_i),$$

$$(I2) \quad -\Delta \left( u_i - \sum_{j \neq i} u_j \right) \geq \lambda_1(D_i)u_i + f_i(u_i) - \sum_{j \neq i} (\lambda_1(D_j)u_j + f_j(u_j)).$$

**Proof**

For a fixed index  $i$ , let us introduce

$$\widehat{u}_i = u_i - \sum_{j \neq i} u_j.$$

Let  $\varphi \geq 0$ ,  $\varphi \in W_0^1(\Omega)$ , and, for  $t > 0$  very small, let us define a new test function  $V = (v_1, \dots, v_k)$ , belonging to  $(W_0^1(\Omega))^k$ , as follows:

$$v_j = \begin{cases} (\widehat{u}_i + t\varphi)^+, & \text{if } j = i, \\ (-u_j + t\varphi)^- = (\widehat{u}_i + t\varphi)^- \chi_{u_j > 0}, & \text{if } j \neq i. \end{cases}$$

We first remark that there is differentiability (with respect to  $t$ ) of all the terms which do not involve derivatives. Indeed, since the map  $u \rightarrow (u^+)^r$  is differentiable, we have, for any set of functions  $\eta_j \in L^s(\Omega)$  and  $r > 1$ :

$$\int_{\Omega} \eta_j v_j^r dx = \begin{cases} \int_{\Omega} \eta_j u_j^r dx + rt \int_{\Omega} \eta_j u_j^{r-1} \varphi dx + o(t), & \text{if } j = i, \\ \int_{\Omega} \eta_j u_j^r dx - rt \int_{\Omega} \eta_j u_j^{r-1} \varphi dx + o(t), & \text{if } j \neq i. \end{cases}$$

By the Sobolev Embedding Theorem, this expansion holds with respect to the  $W_0^1(\Omega)$ -norm provided  $s \in (1, +\infty]$  and  $r \leq (1 - 1/s)(2N/(N - 2))$ . As a first application, letting

$$\alpha_j = \frac{1}{t} \left\{ \int_{\Omega} |v_j|^2 dx - \int_{\Omega} |u_j|^2 dx \right\},$$

and  $r = 2$ , we have

$$\alpha_j = \begin{cases} 2 \int_{\Omega} u_j \varphi dx + o(1), & \text{if } j = i, \\ -2 \int_{\Omega} u_j \varphi dx + o(1), & \text{if } j \neq i. \end{cases}$$

Moreover, letting

$$\beta_j = \frac{1}{t} \left\{ \left( 1 - \frac{\int_{\Omega} v_j(x)^q \widetilde{\phi}_j(x)^q dx}{\left( \int_{\Omega} v_j(x)^{2q} dx \int_{\Omega} \widetilde{\phi}_j(x)^{2q} dx \right)^{1/2}} \right) - \left( 1 - \frac{\int_{\Omega} u_j(x)^q \widetilde{\phi}_j(x)^q dx}{\left( \int_{\Omega} u_j(x)^{2q} dx \int_{\Omega} \widetilde{\phi}_j(x)^{2q} dx \right)^{1/2}} \right) \right\},$$

we find, recalling that  $q \in (1, N/(N-2))$ , by the usual differentiation rules

$$\beta_j = \begin{cases} \frac{-q}{\left(\int_{\Omega} u_j(x)^{2q} dx \int_{\Omega} \tilde{\phi}_j(x)^{2q} dx\right)^{1/2}} \left[ \int_{\Omega} u_j(x)^{q-1} \tilde{\phi}_j(x)^q \varphi(x) dx - \frac{\int_{\Omega} u_j(x)^{2q-1} \varphi(x) dx \int_{\Omega} u_j(x)^q \tilde{\phi}_j(x)^q dx}{\left(\int_{\Omega} u_j(x)^{2q} dx\right)} \right] \\ +o(1), \quad \text{if } j = i, \\ \\ \frac{+q}{\left(\int_{\Omega} u_j(x)^{2q} dx \int_{\Omega} \tilde{\phi}_j(x)^{2q} dx\right)^{1/2}} \left[ \int_{\Omega} u_j(x)^{q-1} \tilde{\phi}_j(x)^q \varphi(x) dx - \frac{\int_{\Omega} u_j(x)^{2q-1} \varphi(x) dx \int_{\Omega} u_j(x)^q \tilde{\phi}_j(x)^q dx}{\left(\int_{\Omega} u_j(x)^{2q} dx\right)} \right] \\ +o(1), \quad \text{if } j \neq i. \end{cases}$$

On the other hand, differentiation with respect to  $t$  may fail when we consider the gradient integrals. Let us denote

$$\delta_j = \frac{1}{t} \left\{ \int_{\Omega} |\nabla v_j|^2 dx - \int_{\Omega} |\nabla u_j|^2 dx \right\}.$$

Although  $t\delta_j \rightarrow 0$  as  $t \rightarrow 0$ , the  $\delta_j$ 's themselves can be unbounded, in general, for they involve boundary integrals which are not necessarily finite for functions in  $W_0^1(\Omega)$ . On the other hand, from the definition

$$\int_{\Omega} |\nabla v_j|^2 dx \leq \int_{\Omega} |\nabla(u_j - t\varphi)|^2 dx, \quad \text{if } j \neq i,$$

we can easily deduce that

$$\delta_j \leq -2 \int_{\Omega} \nabla u_j \cdot \nabla \varphi dx + o(1), \quad \text{if } j \neq i, \text{ and } \varphi \geq 0, \quad (3.10)$$

while, from

$$\begin{aligned} t \sum_j \delta_j &= \sum_j \left\{ \int_{\Omega} |\nabla v_j|^2 dx - \int_{\Omega} |\nabla u_j|^2 dx \right\} \\ &= \int_{\Omega} |\nabla(\hat{u}_i + t\varphi)|^2 dx - \int_{\hat{u}_i + t\varphi=0} |\nabla \hat{u}_i + t\varphi|^2 dx - \int_{\Omega} |\nabla \hat{u}_i|^2 dx + \int_{\hat{u}_i=0} |\nabla \hat{u}_i|^2 dx \\ &= 2t \int_{\Omega} \nabla \hat{u}_i \cdot \nabla \varphi dx + t^2 \int_{\Omega} |\nabla \varphi|^2 dx, \end{aligned}$$

we easily conclude that

$$\sum_j \delta_j = 2 \int_{\Omega} \nabla \hat{u}_i \cdot \nabla \varphi dx + o(1). \quad (3.11)$$

Let us estimate, for a fixed index  $j$ , the difference:

$$\left( \frac{\int_{\Omega} |\nabla v_j(x)|^2 dx}{\int_{\Omega} |v_j(x)|^2 dx} \right)^p - \left( \frac{\int_{\Omega} |\nabla u_j(x)|^2 dx}{\int_{\Omega} |u_j(x)|^2 dx} \right)^p = pt\Lambda^p (\delta_j - \lambda_1(D_j)\alpha_j + o(\delta_j)),$$

here we used the normalization condition (3.7), which implies

$$\int_{\Omega} |\nabla u_j|^2 dx = \left( \frac{\lambda_1(D_j)}{\Lambda} \right)^p. \quad (3.12)$$

On the other hand, we have:

$$\left( 1 - \frac{\int_{\Omega} v_j(x)^q \tilde{\phi}_j(x)^q dx}{\left( \int_{\Omega} v_j(x)^{2q} dx \int_{\Omega} \tilde{\phi}_j(x)^{2q} dx \right)^{1/2}} \right) - \left( 1 - \frac{\int_{\Omega} u_j(x)^q \tilde{\phi}_j(x)^q dx}{\left( \int_{\Omega} u_j(x)^{2q} dx \int_{\Omega} \tilde{\phi}_j(x)^{2q} dx \right)^{1/2}} \right) = t\beta_j.$$

Now we prove inequality (I1). We select  $j \neq i$  and we replace only the  $j$ 'th component  $u_j$  with  $v_j$ . We obtain, as  $t \rightarrow 0^+$ ,

$$\begin{aligned} 0 &\leq \frac{1}{t} (\mathcal{F}_{k,p}(u_1, \dots, v_j, \dots, u_k) - \mathcal{F}_{k,p}(u_1, \dots, u_j, \dots, u_k)) \\ &= \frac{1}{\gamma} (\delta_j - \lambda_1(D_j)\alpha_j - \gamma\beta_j) + o(\delta_j) + o(1) \end{aligned}$$

This inequality, and the boundedness of the  $\alpha_j$ 's, the  $\beta_j$ 's and  $\gamma$  gives a lower bound of the  $\delta_j$ 's. On the other hand (3.10) gives an upper bound of the  $\delta_j$ 's, which are consequently bounded as  $t \rightarrow 0$ . Hence we can deduce from (3.10) and the last inequality that

$$0 \leq -2 \int_{\Omega} (\nabla u_j \cdot \nabla \varphi - \lambda_1(D_j)u_j\varphi - f_j(u_j)\varphi) dx.$$

Since this holds for every pair of indices  $i \neq j$  (though here  $i$  does not appear) and every non negative test function  $\varphi$ , inequality (I1) is proved.

To prove inequality (I2), we argue by contradiction and we assume the existence of  $\varphi \geq 0$  such that

$$\int_{\Omega} \nabla \hat{u}_i \cdot \nabla \varphi dx < \int_{\Omega} \left( \lambda_1(D_i)u_i(x) + f_i(u_i)(x) - \sum_{j \neq i} \lambda_1(D_j)u_j(x) + f_j(u_j)(x) \right) \varphi(x) dx,$$

or, in other words,

$$\int_{\Omega} \nabla \hat{u}_i \cdot \nabla \varphi dx < \sum_i \left( \lambda_1(D_j) \frac{\alpha_j}{2} + \gamma \frac{\beta_j}{2} \right) + o(1). \quad (3.13)$$

Now, by the minimization property of  $U$ , we have,

$$0 \leq \mathcal{F}_{k,p}(v_1, \dots, v_k) - \mathcal{F}_{k,p}(u_1, \dots, u_k) = \frac{t}{\gamma} \sum_j (\delta_j - \lambda_1(D_j)\alpha_j - \gamma\beta_j) + o(t),$$

in contradiction with (3.13) and (3.11).

Theorem 3.4 will easily follow from the next two results:

**Lemma 3.12**

As  $p \rightarrow +\infty$  any family of minimizers of (3.4) satisfying the normalization condition (3.7) with  $\Lambda = \mathfrak{L}_k$  converges, up to a subsequence, to a multiple  $(a_1 \tilde{\phi}_1, \dots, a_k \tilde{\phi}_k)$  ( $a_i \geq 0$ , not all vanishing) strongly in  $(W_0^1(\Omega))^k$ .

**Proof**

From Proposition 3.9 we have, for the minimizers  $u_{i,p}$  and the corresponding  $D_{i,p}$  (we now mention the reference to  $p$  which will then tend to  $+\infty$ ),

$$\frac{1}{k} \sum_i \int_{\Omega} |\nabla u_{i,p}|^2 dx = \frac{1}{k} \sum_i \left( \frac{\lambda_1(D_{i,p})}{\Lambda} \right)^p \leq \left( \frac{\mathcal{M}_{k,p}}{\mathfrak{L}_k} \right)^p \leq 1,$$

while

$$\frac{1}{k} \sum_i \int_{\Omega} |\nabla u_{i,p}|^2 dx \geq \left( \frac{\mathfrak{L}_{k,p}}{\mathfrak{L}_k} \right)^p \geq \frac{1}{k}.$$

Hence the family is bounded in  $(W_0^1(\Omega))^k$  and does not vanish. We extract a sequence  $(u_{i,p_n})_{n \in \mathbb{N}}$  possessing a limit, in the weak  $(W_0^1(\Omega))^k$ -topology and in any  $(L^r(\Omega))^k$ , for subcritical  $r$ 's. We denote by  $(\tilde{u}_i)_{i=1, \dots, k}$  this limit. We infer that the weak limit cannot be identically zero. We have indeed:

$$\lambda_1(D_{i,p_n}) = \frac{\int_{\Omega} |\nabla u_{i,p_n}|^2 dx}{\int_{\Omega} |u_{i,p_n}|^2 dx} \leq k^{1/p_n} \mathfrak{L}_k, \quad \forall i = 1, \dots, k,$$

so that

$$\frac{1}{k} \sum_i \int_{\Omega} |u_{i,p_n}|^2 dx \geq \frac{1}{k^{1/p_n} \mathfrak{L}_k} \sum_i \int_{\Omega} |\nabla u_{i,p_n}|^2 dx.$$

We further remark that, if for some  $i$  the weak limit happens to be zero, then the strong limit vanishes too.

We claim that, for suitable non negative  $a_i$ 's we have  $\tilde{u}_i = a_i \tilde{\phi}_i$ . This is obvious if  $\tilde{u}_i \equiv 0$ . If not, since by Proposition 3.9

$$\lim_{p \rightarrow +\infty} (\mathcal{M}_{k,p} - \mathfrak{L}_{k,p}) = 0,$$

we deduce that, whenever  $\tilde{u}_i \not\equiv 0$ ,

$$1 - \frac{\int_{\Omega} \tilde{u}_i(x)^q \tilde{\phi}_i(x)^q dx}{\left( \int_{\Omega} \tilde{u}_i(x)^{2q} dx \int_{\Omega} \tilde{\phi}_i(x)^{2q} dx \right)^{1/2}} = 0,$$

and therefore that  $\tilde{u}_i$  is a multiple of  $\tilde{\phi}_i$ . To pass from weak to strong convergence we first notice that each  $u_{i,p_n} - \tilde{u}_i$  converges weakly and strongly in  $L^2$  to zero. Now we recall that  $u_{i,p_n}$  satisfy inequalities (I1)–(I2) of Lemma 3.11. Let us multiply (I1) by  $(u_{i,p_n} - \tilde{u}_i)^+$ , (I2) by  $(u_{i,p_n} - \tilde{u}_i)^-$  and take the difference. We obtain

$$\int_{\Omega} \nabla u_{i,p_n} \cdot \nabla (u_{i,p_n} - \tilde{u}_i) dx \leq \int_{\Omega} \sum_{j \neq i} \nabla u_{j,p_n} \cdot \nabla (u_{i,p_n} - \tilde{u}_i)^- dx + o(1).$$

Since  $u_{i,p_n}(x)u_{j,p_n}(x)$  vanishes almost everywhere and  $\tilde{u}_i \geq 0$ , we infer

$$\begin{aligned} \int_{\Omega} \nabla u_{i,p_n} \cdot \nabla (u_{i,p_n} - \tilde{u}_i) dx &\leq \sum_{j \neq i} \int_{\Omega} \nabla u_{j,p_n} \cdot \nabla \tilde{u}_i dx + o(1) \\ &= \sum_{j \neq i} \int_{\Omega} \nabla \tilde{u}_j \cdot \nabla \tilde{u}_i dx + o(1) = o(1). \end{aligned}$$

Thus we can deduce strong convergence from the weak. We have indeed

$$\|\nabla(u_{i,p_n} - \tilde{u}_i)\|^2 = \int_{\Omega} \nabla u_{i,p_n} \cdot \nabla (u_{i,p_n} - \tilde{u}_i) dx - \int_{\Omega} \nabla \tilde{u}_i \cdot \nabla (u_{i,p_n} - \tilde{u}_i) dx = o(1).$$

### Lemma 3.13

Let  $U_n = (u_{1,p_n}, \dots, u_{k,p_n})$  ( $n \in \mathbb{N}$ ) as in the proof of Lemma 3.12. Then its limit, as  $n \rightarrow +\infty$ ,  $\tilde{U} := (\tilde{u}_1, \dots, \tilde{u}_k)$  verifies the inequalities in the statement of Theorem 3.4.

### Proof

First of all, we wish to pass to the limit in formulas (I1)–(I2) of Lemma 3.11, in the sense of distributions. From the previous lemma we deduce that  $-\Delta u_{i,p_n} \rightarrow -\Delta \tilde{u}_i$  in  $H^{-1}(\Omega)$ . Hence one can pass to the limit in inequality (I1). Let us turn to (I2). We remark that also  $f_i(u_{i,p_n})$  converge to  $f_i(\tilde{u}_i)$ , provided  $\tilde{u}_i \neq 0$ . For such  $i$ 's inequality (I2) passes to the limit, because so does its right hand. On the other hand, when the limit  $\tilde{u}_i$  does vanish then (I2) holds because of (I1) and the fact that  $-\Delta \tilde{u}_i = 0$ . In order to end the proof, we have to prove convergence of the eigenvalues  $\lambda_1(D_{i,p_n})$  to  $\mathfrak{L}_k$  whenever  $\tilde{u}_i$  does not vanish identically. At first we notice that the  $\lambda_1(D_{i,p_n})$ 's do converge, thanks to the strong convergence of the  $u_{i,p_n}$ 's to limits  $\lambda_1(\tilde{D}_i) \leq \mathfrak{L}_k$ . Assuming  $\lambda_1(\tilde{D}_i) < \mathfrak{L}_k$  we deduce from (3.12), using again the strong convergence, that  $\tilde{u}_i \equiv 0$ .

### Remark 3.14

(a) Thanks to [8], Theorem 8.3 all the  $u_{i,p}$ 's and their limits  $\tilde{u}_i$  are locally Lipschitz continuous in the interior of  $\Omega$  and continuous up to the

boundary, for (I1), if the boundary  $\partial\Omega$  is Lipschitz, or has the interior cone property; moreover, they are globally Lipschitz up to the boundary, if  $\partial\Omega$  is of class  $\mathcal{C}^1$ .

(b) Of course, since the eigenfunctions are normalized in  $L^2$

$$a_i \stackrel{\text{def}}{=} \|\tilde{u}_i\|_2.$$

In general, it may happen that some of the  $a_i$ 's vanish; we denote

$$\mathbf{k}_0 = \{i \in \{1, \dots, k\} : a_i = 0\}. \quad (3.14)$$

(c) Going back to the proof of Lemma 3.12, we can extract a subsequence with the further property that the  $u_{i,p_n}/\|u_{i,p_n}\|_2$  converge strongly in  $L^2(\Omega)$  and weakly in  $W_0^1(\Omega)$  to  $\tilde{\phi}_i$ , also for those indices  $i$ 's for which the component  $u_{i,p_n}$  normalized as in (3.12) strongly converges to 0.

(d) We also infer from (3.12) that

$$\lambda_1(\tilde{D}_i) \leq \mathfrak{L}_k \quad \forall i \in \{1, \dots, k\},$$

while

$$\lambda_1(\tilde{D}_i) = \mathfrak{L}_k, \quad \forall i \notin \mathbf{k}_0.$$

(e) Obviously the system of differential inequalities (I1)-(I2) are fulfilled by the set  $(\tilde{u}_i)$  with  $i \notin \mathbf{k}_0$ .

(f) For  $i \notin \mathbf{k}_0$ , the  $\tilde{D}_i$ 's are open and possess finitely many connected components.

## 4 Further results in two dimensions

In this section, we recall from the introduction that we work under Assumptions 1.1 and 1.2. This implies that  $\partial\Omega$  has finitely many connected components. For simplicity, we shall omit the potential  $V$  in the following discussion. All the arguments can be straightforwardly extended (sometimes at the price to replace  $\mathcal{C}^2$  by  $\mathcal{C}^{1,1^-} := \cup_{\alpha < 1} \mathcal{C}^{1,\alpha}$ ) in order to cover the case of a non vanishing bounded potential. A special caution is only due in the proof of Theorem 4.6, where the needed extra argument is outlined.

## 4.1 Case when $\mathbf{k}_0 = \emptyset$ .

In this section we discard in a first step all the identically vanishing components. Hence, from now on, we will assume that

$$\mathbf{k}_0 = \emptyset .$$

Re-labeling if necessary, and taking a smaller  $k$ , we assume that the components of

$$U = (u_1, \dots, u_k)$$

are non negative, non vanishing  $W_0^1(\Omega)$ -functions, such that  $u_i(x)u_j(x) \equiv 0$  almost everywhere in  $\Omega$  (for  $i \neq j$ ), satisfying the two differential inequalities (I1)-(I2) of Theorem 3.4. As a consequence of Remark 3.14 (a), they are continuous on the closure of  $\Omega$ . Hence, by expanding the set of indices if necessary, we can always assume that the sets

$$D_i = \{u_i > 0\}$$

are open and connected.

Let us define the set of zeroes of  $U$  as

$$\mathcal{Z} = \{x \in \Omega : u_i(x) = 0, \forall i = 1, \dots, k\},$$

and define the *multiplicity* of  $x \in \Omega$  as the number  $m(x)$ :

$$m(x) = \#\{i : \text{meas}(\{u_i > 0\} \cap B_r(x)) > 0, \forall r > 0\}. \quad (4.1)$$

We shall denote by

$$\mathcal{Z}_h = \{x \in \Omega : m(x) \geq h\}$$

the set of points of multiplicity greater than or equal to the integer  $h$  and by

$$\mathcal{Z}^h = \{x \in \Omega : m(x) = h\}.$$

We remark that, by definition,  $\mathcal{Z}^0$  is open. Let us now consider  $\mathcal{Z}^1$ . This is the object of:

### Proposition 4.1

$$\mathcal{Z}^1 = \bigcup_{i=1}^k D_i .$$

**Proof**

If a ball  $B_r(x_0)$  intersects only  $D_i$  we deduce from (I1)–(I2) that  $u_i$  is a non negative solution to the differential equation  $-\Delta u = \Lambda u$  on  $B_r(x_0)$ . By the strong maximum principle then  $u_i$  is strictly positive on  $B_r(x_0)$  and therefore  $B_r(x_0) \subset D_i$  and all the points of  $B_r(x_0)$  have multiplicity equal to one. ■

To continue the analysis of the topological properties related to points of multiplicity two and more, we shall consider the simplicial homology groups with coefficients in  $\mathbb{Z}_2$ ,  $H_n(X)$ , for  $n = 0, 1$ . We recall that  $\text{rank}(H_0(X))$  is the number of connected components of  $X$ . For open subsets of an euclidean space, as the fundamental group  $\pi_1$  is already abelian, there holds  $\text{rank}(\pi_1(X)) = \text{rank}(H_1(X))$ . Finally for planar bounded open subsets,  $\text{rank}(H_1(X)) + 1$  is the number of connected components of  $\partial X$ . A reference book for the algebraic topology concepts is the Greenberg and Harper's book [13].

**Proposition 4.2**

*If  $H_1(\Omega)$  is finite, so is  $H_1(D_i)$ , for every  $i$ .*

**Proof**

Let us consider a loop  $\gamma \subset D_i$  which is homotopically trivial in  $\Omega$  but not in  $D_i$ ; hence denoting by  $\Sigma$  the inner region of  $\gamma$ , we have that  $\Sigma \subset \Omega$  but  $\Sigma \not\subset D_i$ . Let  $j \neq i$ ; then either  $D_j$  is contained in  $\Sigma$  or in its complement, for it is connected. To prove the proposition, we argue by contradiction and we assume that  $H_1(D_i)$  is infinite. Then also  $\text{rank}(\pi_1(D_i))$  is infinite; thus we infer the existence of at least one loop  $\gamma = \partial\Sigma$  such that  $\Sigma \cap \bigcup_{j \neq i} D_j = \emptyset$  and  $\Sigma \not\subset D_i$ . Since all the  $u_j$ 's ( $j \neq i$ ) vanish identically in  $\Sigma$ , we deduce from (I1)–(I2) that  $-\Delta u_i = \Lambda u_i$  in  $\Sigma$  and, by the strong maximum principle, that  $u_i$  is strictly positive there; thus  $\Sigma \subset D_i$ , a contradiction. ■

Let us consider

$$\Gamma_{i,j} = \partial D_i \cap \partial D_j \cap \mathcal{Z}^2,$$

$$D_{i,j} = D_i \cup D_j \cup \Gamma_{i,j}.$$

Our next goal consists in showing that the  $\Gamma_{i,j}$  consist in a finite number of (possibly open) arcs. This will require some topological considerations.

**Proposition 4.3**

*Let  $x_0 \in \Omega$  such that  $m(x_0) = 2$ . Then  $u_i - u_j$  is in  $\mathcal{C}^{1,1-}$  in some neighborhood of  $x_0$  and  $\nabla(u_i - u_j)(x_0) \neq 0$ . Furthermore  $\mathcal{Z}^2$  is locally a  $\mathcal{C}^{1,+}$ -curve through  $x_0$ .*

**Proof**

Relabeling, we can always assume  $x_0 \in \partial\{u_1 > 0\} \cap \partial\{u_2 > 0\}$ ; thus for all  $r$  small enough  $B(x_0, r) \cap D_i = \emptyset$  for all  $i > 2$ . Then  $u = u_1 - u_2$  satisfies the equation  $-\Delta u = \Lambda u$  in  $B(x_0, r)$  and is consequently  $\mathcal{C}^{1,1-}$  near  $x_0$ . Therefore the zero set of  $u$  near  $x_0$  (by a standard result on the zero set) is made up by a finite number of even regular curves starting from  $x_0$ . But there are actually only two arcs meeting at  $x_0$ . Indeed, if not, at least one of the  $D_i$ 's should be disconnected. In this case the zero set is actually locally a regular line passing through  $x_0$  and the Boundary Point Lemma gives the proposition. ■

**Proposition 4.4**

*If  $H_1(\Omega)$  is finite so is  $H_1(D_{i,j})$ , for every  $i \neq j$ .*

**Proof**

This is an obvious statement if  $\Gamma_{i,j} = \emptyset$ .

Let us consider a loop  $\gamma \subset D_{i,j}$  which is trivial in  $\Omega$  but not in  $D_{i,j}$ ; hence  $\gamma = \partial\Sigma$  with  $\Sigma \subset \Omega$  but  $\Sigma \not\subset D_{i,j}$ . Let  $\ell \notin \{i, j\}$ ; then either  $D_\ell$  is contained in  $\Sigma$  or in its complement. Thus, arguing by contradiction we infer the existence of a loop  $\gamma = \partial\Sigma$  such that  $\Sigma \cap \bigcup_{\ell \neq i,j} D_\ell = \emptyset$  and  $\Sigma \not\subset D_{i,j}$ . Since all the  $u_\ell$ 's ( $\ell \neq i, j$ ) vanish identically in  $\Sigma$ , we deduce from (I1)-(I2) that  $-\Delta(u_i - u_j) = \Lambda(u_i - u_j)$  in  $\Sigma$  and, from Proposition 4.3 and the unique continuation principle we infer that  $\mathcal{Z} \cap \Sigma \subset \Gamma_{i,j}$ ; thus  $\Sigma \subset D_{i,j}$ , a contradiction. ■

**Lemma 4.5**

*If  $H_1(D_{i,j})$  is finite, then  $\Gamma_{i,j}$  has finitely many connected components.*

**Proof**

To prove the statement we will take advantage of the Mayer–Vietoris Theorem. The Mayer–Vietoris sequence is usually proven to be exact for a triad  $X, X_1$  and  $X_2$  where  $X_1$  and  $X_2$  are open subsets of the topological space  $X$  and  $X$  is the union of  $X_1$  and  $X_2$  (such a triplet is called an admissible triad). Here we would like to apply the Mayer–Vietoris sequence to the triad  $X = D_{i,j}$ ,  $X_1 = D_i \cup \Gamma_{i,j}$  and  $X_2 = D_j \cup \Gamma_{i,j}$ , but these latter two are not open in  $D_{i,j}$ . However, the Mayer–Vietoris sequence is still available because, thanks to Proposition 4.3, the  $\Gamma_{i,j}$ 's are regular embedded one-dimensional submanifolds in  $D_{i,j}$ . Hence each  $D_\ell \cup \Gamma_{i,j}$  is a euclidean neighborhood retract in  $D_{i,j}$  and, as such it has the same homology as the corresponding  $D_\ell$  ( $\ell = i, j$ ). Thus, following [13], the triplet  $D_{i,j}, D_i \cup \Gamma_{i,j}, D_j \cup \Gamma_{i,j}$  is a proper

excision triad and thus the Mayer–Vietoris sequence is exact:

$$H_1(D_{i,j}) \xrightarrow{\partial_*} H_0(\Gamma_{i,j}) \xrightarrow{i_* \oplus -j_*} H_0(D_i \cup \Gamma_{i,j}) \oplus H_0(D_j \cup \Gamma_{i,j})$$

The assertion then follows as a consequence of Propositions 4.2 and 4.4 taking into account of the connectedness of  $D_\ell$  ( $\ell = i, j$ ). Indeed since both  $H_1(D_{i,j})$  and  $H_0(D_i \cup \Gamma_{i,j}) \oplus H_0(D_j \cup \Gamma_{i,j}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$  are finite, we have that both the range of  $\partial_*$  and the coker of  $i_* \oplus -j_*$  are finite and thus so is  $H_0(\Gamma_{i,j})$ . ■

The following results follow from [8]:

**Theorem 4.6**

We have,  $\forall i = 1, \dots, k$  :

(a)  $u_i \in W_{loc}^{1,\infty}(\Omega)$ .

(b) The function  $\sum_{i=1}^k |\nabla u_i|$  admits a continuous representative in  $\Omega$ .

(c) If  $x_0 \in \mathcal{Z}_3$  then,

$$\lim_{x \rightarrow x_0} \sum_{i=1}^k |\nabla u_i(x)| = 0 .$$

(d) If  $x_0 \in \mathcal{Z}_3$  then,

$$\lim_{x \rightarrow x_0} \frac{u_i(x)}{|x - x_0|} = 0 .$$

**Proof**

Part (a) is indeed stated as Theorem 8.3 of [8]. To prove part (b), we first observe that locally in  $\bigcup_{i=0}^2 \mathcal{Z}^i$  (which is open in  $\Omega$ ) the  $u_i$ 's satisfy the differential equation  $-\Delta(u_i - u_j) = \Lambda(u_i - u_j)$ : hence they are regular (of class  $\mathcal{C}^1(\bigcup_{i=0}^2 \mathcal{Z}^i)$ ). The continuity up to the  $\mathcal{Z}_3$  (which is indeed the boundary of  $\bigcup_{i=0}^2 \mathcal{Z}^i$ ) is then a consequence of the vanishing of the limit stated in part (c), which is indeed Theorem 9.3 in [8]. The proof of Theorem 9.3 was originally performed in the absence of the  $L^\infty$  potential, but all the arguments can be promptly adapted to cover also this case. A special care is needed when, at the beginning, the function  $\sum_{i=1}^k |\nabla u_i|^2$  is shown to be a subsolution to a linear differential equation. Subsequently the mean value property is applied. This is not exactly true in the present situation, for the linear problem is now perturbed by a term of the form  $\nabla w \cdot \nabla((\Lambda + V(x))w)$ , where  $w$  is an auxiliary function (which can be taken, by the way, the same as defined in (4.3)). After integrating by parts one easily sees that the contribution of this

term is negligible. This observation permits to follow, from then on, exactly the same arguments of the quoted paper.

**Lemma 4.7**

Let  $x_0 \in \mathcal{Z}_3$ . Then there exists a sequence  $\{x_n\} \subset \Omega$  such that  $m(x_n) = 2$  and  $x_n \rightarrow x_0$  as  $n \rightarrow +\infty$ .

**Proof**

Assume not, then there would be an element  $y_0$  of  $\mathcal{Z}_3$  having a positive distance  $d$  from  $\mathcal{Z}^2$ . Let  $r < d/2$ ; then the ball  $B(y_0, r)$  intersects at least three of the  $D_i$ 's; therefore there exist  $i \in \{1, \dots, k\}$ ,  $x \in D_i$  and  $z_0 \in \mathcal{Z}_3$  such that  $\rho = d(x, z_0) = d(x, \mathcal{Z}_3) < d(x, \mathcal{Z}^2)$ . Then the ball  $B(x, \rho)$  is tangent from the interior of  $D_i$  to  $\mathcal{Z}_3$  in  $z_0$ . Furthermore  $u_i$  solves an elliptic PDE (in the sense of [12]) and it is positive on  $D_i$ . Then we thus infer from the Boundary Point Lemma (see Lemma 3.4 and formula (3.11) in [12] ) that

$$\liminf_{h \rightarrow 0} \frac{u_i(z_0 + h\nu)}{h} > 0, \tag{4.2}$$

where  $\nu$  denotes the inner normal to  $\partial B(x, \rho)$  at  $z_0$ , in contrast with Theorem 4.6(c). ■

A simple but important consequence of this discussion is the following result (the closure is taken in  $\overline{\Omega}$ ):

**Proposition 4.8**

If  $H_1(\Omega)$  is finite then  $\overline{\mathcal{Z}_3}$  has a finite number of connected components.

**Proof**

Indeed, we have  $\overline{\mathcal{Z}_3} \subset \cup_{i,j} (\overline{\Gamma_{i,j}} \setminus \Gamma_{i,j})$ , and each  $\Gamma_{i,j}$  is the union of finitely many arcs, each of them being homeomorphic to the real line (non compact case) or to a circle. In the first case they have connected and closed  $\alpha$  and  $\omega$ -limits.

We recall that the  $\alpha$  and  $\omega$ -limits of a parametrized arc  $\Gamma(t)$  are the sets of the limit points as the parameter tends to  $-\infty$  and  $+\infty$  respectively: one easily sees that bounded arcs have compact and connected  $\alpha$  and  $\omega$ -limits. Therefore each  $\overline{\Gamma_{i,j}} \setminus \Gamma_{i,j}$  contributes with finitely many connected components by the previous proposition. ■

In addition we have:

**Proposition 4.9**

If  $H_1(\Omega)$  is finite then  $\overline{\mathcal{Z}_3 \cup \mathcal{Z}^0}$  has a finite number of connected components.

**Proof**

Indeed, by Theorem 4.6 and Lemma 4.7, the boundary  $\partial(\overline{\mathcal{Z}_3 \cup \mathcal{Z}^0})$  is the same as  $\partial\overline{\mathcal{Z}_3}$  and this last one has finitely many connected components.

Our next goal is the following

**Theorem 4.10**

An isolated connected component of  $\overline{\mathcal{Z}_3 \cup \mathcal{Z}^0}$  consists of a single point.

This theorem implies straightforwardly, having in mind that  $\mathcal{Z}^0$  is open, that

**Corollary 4.11**

If  $H_1(\Omega)$  is finite then  $\mathcal{Z}^0 = \emptyset$ .

**Proof of Theorem 4.10**

To prove the theorem we focus on a connected component  $Y_0$  of  $\overline{\mathcal{Z}_3 \cup \mathcal{Z}^0}$  and we show that it is reduced to a single point. First we consider an open and connected neighbourhood  $\mathcal{N}$  of  $Y_0$  in  $\mathbb{R}^2$  having a regular boundary and such that  $\overline{\mathcal{N} \cap \overline{\mathcal{Z}_3 \cup \mathcal{Z}^0}} = Y_0$ . Since  $Y_0$  is connected, we can choose  $\mathcal{N}$  in such a way that its boundary has exactly one or two connected components (each diffeomorphic to  $S^1$ ), depending on whether  $Y_0$  disconnects  $\mathbb{R}^2$  or not. This can be achieved by taking one connected component of a regular sublevel of a non negative  $C^\infty$  function having  $Y_0$  as null set. Furthermore we can take the measure of  $\mathcal{N} \setminus Y_0$  small enough that none of the  $D_i$ 's is completely enclosed in  $\mathcal{N}$ . We may assume that  $\partial\mathcal{N}$  intersects transversally the  $\Gamma_{i,j}$ 's and  $\partial\Omega$ . Thus, after possibly cutting-off some portions of  $\mathcal{N}$ , we can assume that each oriented arc in  $\Gamma_{i,j}$  intersects  $\partial\mathcal{N}$  exactly once. Then each  $D_i \cap \mathcal{N}$  has a finite number of connected components. Moreover, we can manage to have these intersections simply connected. Indeed, assuming not, and arguing as in the proof of Proposition 4.2, at least one of the other  $D_j$ 's should be entirely contained in  $D_i \cap \mathcal{N} \setminus Y_0$ , what we have excluded by taking the measure of  $\mathcal{N} \setminus Y_0$  small enough. We can label the connected components of  $D_i \cap \mathcal{N}$  and  $(\mathbb{R}^2 \setminus \Omega) \cap \mathcal{N}$  clockwise from 1 to  $h$ , according with their intersection with  $\partial\mathcal{N}$ . By taking a double covering of  $\mathcal{N}$ , still denoted by  $\mathcal{N}$ , if necessary, we may assume that  $h$  is an even integer. Indeed, assuming  $h$  to be odd, we choose some point  $y_0 \in Y_0$  and perform a double covering branched at  $y_0$  (in complex notation  $f(z) = (z - y_0)^2$ ). With some abuse of notation, we denote with the same symbols  $D_i$ ,  $\mathcal{N}$  and  $Y_0$  and their pre-images, and we denote

$\tilde{u}_i(z) = u_i(f(z))$ . Note that

$$\Delta \tilde{u}_i(z) = 8|z - y_0|^2 \Delta u_i(f(z)) .$$

If the original  $h$  was even we set  $\tilde{u}_i = u_i$  and we define, in both cases, the auxiliary function

$$w(x) = \begin{cases} \sigma(x)\tilde{u}_i(x) , & \text{if } x \in D_i \\ 0 & \text{otherwise} \end{cases} \quad (4.3)$$

where  $\sigma$  is a sign assignment compatible with the partition of  $\mathcal{N}$ . The function  $w$  satisfies the linear equation:

$$-\Delta w = \Lambda a(x)w \quad \text{in } \Omega \cap (\mathcal{N} \setminus Y_0) \quad (4.4)$$

$$w = 0 \quad \text{on } (\partial\Omega \cap \mathcal{N}) \cup Y_0 , \quad (4.5)$$

where  $a \equiv 1$  if no double covering has been performed, otherwise  $a(x) = 8|x - y_0|^2$  is the conformal factor. Of course  $w$  is regular outside the singular component  $Y_0$ . Also,  $w$  vanishes identically on the open set  $N \cap \mathcal{Z}^0$ ; hence, thanks to Theorem 4.6 (c),  $w$  is in  $\mathcal{C}^1(\Omega \cap \mathcal{N})$

We claim that  $w$  actually solves (4.4) in the whole of  $\Omega \cap \mathcal{N}$ . This is not a trivial fact, for there is no a priori bound on the Hausdorff measure of the boundary  $\partial Y_0$  of the singular set. In order to overcome this problem we make the following construction:

**Proposition 4.12**

*There exists a family of neighborhoods  $\mathcal{N}_\delta \subset \mathcal{N}$ , decreasing to  $Y_0$  and bounded by a finite number of regular arcs, with the property that*

$$\lim_{\delta \rightarrow 0} \int_{\partial(\mathcal{N}_\delta \cap \Omega)} |\nabla w| ds = 0 , \quad (4.6)$$

where  $ds$  denotes the measure on the union of these regular arcs.

Postponing the proof of the proposition, we end the proof of Theorem 4.10. Testing (4.4) with any function  $\varphi \in C_0^\infty(\mathcal{N} \cap \Omega)$ , it follows by decomposing the integration on  $\mathcal{N} \setminus \mathcal{N}_\delta$  and on  $\mathcal{N}_\delta$  and then making an integration by parts for the second integral:

$$\begin{aligned} & \left| \int_{\mathcal{N} \cap \Omega} (\nabla w \cdot \nabla \varphi - \Lambda a(x)u\varphi) dx \right| \\ &= \left| \int_{\mathcal{N}_\delta \cap \Omega} (\nabla w \nabla \varphi - \Lambda a(x)u\varphi) dx - \int_{\partial(\mathcal{N}_\delta \cap \Omega)} \varphi \nabla w \cdot \nu ds \right| \\ &\leq C(\sup_{\mathcal{N}_\delta \cap \Omega \cap \text{supp}(\varphi)} (|w| + |\nabla w|) + \int_{\partial(\mathcal{N}_\delta \cap \Omega)} |\nabla w| ds) . \end{aligned}$$

We have now to show that the right hand side is  $o(1)$ . Proposition 4.12 ensures that, as  $\delta \rightarrow 0$ ,

$$\int_{\partial(\mathcal{N}_\delta \cap \Omega)} |\nabla w| ds = o(1).$$

So it remains to show that, as  $\delta \rightarrow 0$ , there holds

$$\sup_{\mathcal{N}_\delta \cap \Omega \cap \text{supp}(\varphi)} (|w| + |\nabla w|) = o(1).$$

This can be done by Theorem 4.6. Indeed both  $w$  and  $\nabla w$  are uniformly continuous on  $\text{supp}(\varphi)$  which is compactly contained in  $\Omega$ , and the distance of  $\mathcal{N}_\delta \cap \text{supp}(\varphi)$  to  $Y_0$  tends to zero. Finally we recall that  $w$  and its gradient vanish identically on  $Y_0$ . Hence  $w$  solves (4.4) on the whole of  $\mathcal{N} \cap \Omega$ . By a classical local regularity result by Hartman and Wintner ([15], Corollary 1), we know that interior critical points of solutions to such class of equations are isolated and have finite (local) multiplicity  $m$ , and satisfy (2.1).

Hence we are left with the case when  $Y_0$  is contained in a connected component of the boundary. We notice that, in this case, we do not need the double covering, for a sign assignment compatible with the partition of  $\mathcal{N}$  always exists, since  $\partial\Omega$  disconnects  $\partial\mathcal{N}$ . Now we need an extension of Theorem 2.1, suitable to cover the case of domains possessing a finite number of  $\mathcal{C}^{1,+}$ -corners: namely Theorems 2.3 and 2.6 in Section 2.

In particular, these results guarantee finiteness of critical points also at the boundary.

### Proof of Proposition 4.12

First of all, let us consider a triplet of adjacent domains separated by the two arcs  $\Gamma_{i,j}$  and  $\Gamma_{j,k}$ . We claim that the distance of the arcs, relative to  $D_j$ , must vanish. In other words, we claim that

$$\inf \left\{ \int_0^1 |\dot{\gamma}(s)| ds : \gamma(0) \in \Gamma_{i,j}, \gamma(s) \in D_j, \forall s \in (0,1), \gamma(1) \in \Gamma_{j,k} \right\} = 0.$$

Indeed, if not, one could find in  $D_j$  a ball, tangent to the boundary  $\partial D_j \cap Y_0$  at, say,  $x_0$  and having positive distance from both  $\Gamma_{i,j}$  and  $\Gamma_{j,k}$ . By the Boundary Point Lemma, at  $x_0$  the gradient of  $w$  cannot vanish (in the weak sense of (4.2)), in contradiction with the fact that  $x_0 \in Y_0$ .

As a second remark, by integrating the equation  $-\Delta u_i = \Lambda u_i$  over the set  $\{u_i > \varepsilon\}$  and using the Divergence Theorem, we obtain,

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\partial\{u_i > \varepsilon\}} |\nabla u_i| ds \leq C \int_{\Omega} u_i dx < +\infty.$$

The Divergence Theorem is applicable because the level sets (according to Hartman and Wintner's regularity result) of the eigenfunctions of  $-\Delta + V(x)$  are compact and piecewise  $\mathcal{C}^1$  when the potential  $V$  is bounded. Since the components of  $\partial\{u_i > \varepsilon\}$  converge to  $\Gamma_{i,j}$  as  $\varepsilon \rightarrow 0$ , in  $\mathcal{C}^1$  as parametrized curved, we obtain that

$$\int_{\Gamma_{i,j}} |\nabla w| ds < +\infty, \quad \forall i, j.$$

If the arc  $\Gamma_{i,j}$  meets  $Y_0$ , then we can choose an orientation for its parametrization  $\gamma(t)$  in a way that  $\lim_{t \rightarrow +\infty} \text{dist}(\gamma(t), Y_0) = 0$ , and we have :

$$\begin{aligned} \int_0^{+\infty} |\nabla w(\gamma(t))| |\gamma'(t)| dt &< +\infty; \\ \gamma(t) &\notin Y_0, \quad \forall t \geq 0; \\ \lim_{t \rightarrow +\infty} \text{dist}(\gamma(t), Y_0) &= 0. \end{aligned}$$

Therefore we can conclude

$$\lim_{\delta \rightarrow 0} \int_{\Gamma_{i,j} \cap B_\delta(Y_0)} |\nabla w| ds = 0, \quad \forall i, j,$$

where we denoted  $B_\delta(Y_0) = \{x \in \mathbb{R}^2, d(x, Y_0) < \delta\}$ .

Now we illustrate the construction of the boundary of the neighbourhood  $\mathcal{N}_\delta$ . We start by taking the component of  $\Gamma_{i,j} \cap B_\delta(Y_0)$  ending at  $Y_0$ ; then we can jump from this arc to the next  $\Gamma_{j,k}$ , still remaining in  $B_\delta(Y_0) \cap D_j$ , following a segment of arbitrarily short length. We proceed in this way passing from one arc to the next until we complete the loop. ■

Next result straightforwardly follows from Theorems 2.3 and 2.6 applied to the auxiliary function  $w$  defined in (4.3); it completes our analysis of the asymptotic expansion of the nodal set at multiple intersection points.

#### Theorem 4.13

Let  $x_0 \in \Omega$  with  $m(x_0) = h \geq 3$  that is isolated in  $\overline{\mathcal{Z}}_3$ . Then there exist an integer  $n \geq h$  (the local multiplicity),  $c \in \mathbb{R} \setminus 0$ ,  $\theta_0 \in (-\pi, \pi]$  such that

$$\sum_{i=1}^h u_i(r, \theta) = cr^{\frac{n}{2}} \left| \cos\left(\frac{n}{2}(\theta + \theta_0)\right) \right| + o(r^{\frac{n}{2}}),$$

as  $r \rightarrow 0$ , where  $(r, \theta)$  denotes a system of polar coordinates around  $x_0$  and  $n$  is the local multiplicity of  $x_0$ .

Moreover the nodal set in a neighbourhood of  $x_0$  is the union of  $n$  closed arcs of class  $\mathcal{C}^{1,+}$  meeting at  $x_0$  and spanning angles of opening angle  $2\pi/n$ . If the boundary  $\partial\Omega$  presents a  $\mathcal{C}^{1,+}$ -corner of amplitude  $\alpha\pi$  at  $x_0$  and the nodal set hits  $x_0$  from inside the corner, then there exist an integer  $n$  and  $R > 0$  such that the component of  $\overline{u^{-1}(\{0\})} \cap B(x_0, R)$  lying inside the corner is composed by  $n$   $\mathcal{C}^{1,+}$ -simple arcs which all end in  $x_0$  and whose tangent lines at  $x_0$  divide the sector into  $n + 1$  angles of equal opening angle  $\pi\alpha/(n + 1)$ .

## 4.2 General case.

Let us come back to the general case. We no more assume a priori that  $\mathbf{k}_0 = \emptyset$ , nor the connectedness of the  $D_i$ 's. Then we obtain :

### Theorem 4.14

If  $N = 2$  and  $\Omega$  is bounded, connected, satisfies the interior cone property and with a piecewise  $\mathcal{C}^{1,+}$  boundary, then the assertion of Theorem 3.4 holds with all  $a_i$ 's strictly positive.

Moreover any minimizing partition  $\mathcal{D}$  admits an open regular connected representative.

### Proof

Assume that some of the  $a_i$  vanish or, in other words, that  $\mathbf{k}_0 \neq \emptyset$ . Let us denote

$$\mathcal{W}_0 = \Omega \setminus \bigcup_{i \notin \mathbf{k}_0} D_i$$

the nodal set. Then the 2-dimensional measure of  $\mathcal{W}_0$  is positive, since it contains the supports of  $\tilde{\phi}_i$  for all  $i \in \mathbf{k}_0$ .

Now, as already remarked, inequalities (I1) and (I2) are still available when we discard all the vanishing components and we take  $U = (u_i)_{i \notin \mathbf{k}_0}$ . Hence Theorem 4.13 applies and, as a direct consequence, we find that the zero set  $\mathcal{Z}$ , being the union of a finite number of closed  $\mathcal{C}^{1,+}$  curves has vanishing measure in  $\mathbb{R}^2$ , a contradiction. This also implies that the partition associated with the  $D_i$  is strong and that the zero set is indeed the nodal set  $N(\mathcal{D})$  as defined in (1.9).

Now assume by contradiction some elements of the partition not be connected. We can anyway choose a set of first eigenfunctions having each a connected support, but we clearly have an open, non empty set of multiplicity zero points. This contradicts Corollary 4.11.

**Remark 4.15**

*Note that Theorem 4.14 completes the proof of Theorem 1.12.*

**Proof of Theorem 1.14**

But we also get the proof of Theorem 1.14 in the following way. If the graph associated to  $\tilde{D}$  is bipartite we can find  $\epsilon_i = \pm 1$  satisfying  $\epsilon_i \epsilon_j = -1$  if  $D_i \sim D_j$  and such that  $u := \sum_i \epsilon_i a_i \tilde{\varphi}_i$  is in  $W_0^1(\Omega)$  and satisfies

$$(-\Delta + V)u = \mathfrak{L}_k(\Omega)u \tag{4.7}$$

in  $\Omega \setminus \mathcal{Z}_3$ . But we have proven that  $\mathcal{Z}_3$  consists of isolated points (which cannot be the support of a distribution in  $W^{-1}(\Omega)$ ). Hence (4.7) is satisfied in  $\Omega$  and  $u$  is actually an eigenfunction of  $H(\Omega)$  corresponding to  $\mathfrak{L}_k(\Omega)$ . We refer to Appendix B for a complementary discussion.

## 5 More on nodal sets and partitions

We continue by discussing more deeply the links between the various spectral sequences.

The first important property is given by :

### Proposition 5.1

Let  $H(\Omega)$  be defined as above. Then

$$\mathfrak{L}_k(\Omega) < \mathfrak{L}_{k+1}(\Omega) \text{ for } k \geq 1. \quad (5.1)$$

### Proof

We take indeed a minimal  $(k + 1)$ -partition of  $\Omega$ . We have proved that this partition is regular. If we take any subpartition by  $k$  elements of the previous partitions. This cannot be a  $k$ -minimal partition (it has not the “strong partition” property). So the inequality in (5.1) should be strict.

The second property concerns the **domain monotonicity**. It is indeed immediate to verify :

### Proposition 5.2

If  $\Omega \subset \tilde{\Omega}$ , then

$$\mathfrak{L}_k(\tilde{\Omega}) \leq \mathfrak{L}_k(\Omega), \quad \forall k \geq 1.$$

We observe indeed that each partition of  $\Omega$  is a partition of  $\tilde{\Omega}$ .

### Remark 5.3

The analysis of the equality in the proposition will involve the capacity of  $\tilde{\Omega} \setminus \Omega$ . See [4] and references therein.

We now come back to a definition which was briefly mentioned in the introduction. Having in mind Definition 1.16, we denote, for any integer  $k \geq 1$ , by  $L_k(\Omega)$  the smallest eigenvalue whose eigenspace contains an eigenfunction with  $k$  nodal domains. We take  $L_k = +\infty$ , if there are no eigenfunctions with  $k$  nodal domains. We call this sequence the spectral nodal sequence.

### Proposition 5.4

Let  $\lambda$  be an eigenvalue corresponding to an eigenfunction with  $k$  nodal domains. Then

$$\mathfrak{L}_k \leq \lambda. \quad (5.2)$$

**Proof of Proposition 5.4.**

If  $u$  is an eigenfunction associated with  $\lambda$  and with  $k$  nodal domains, then, taking as  $\mathcal{B}_0$  the collection of these nodal domains, we obtain :

$$\inf_{\mathcal{B} \in \mathfrak{D}_k} \Lambda(\mathcal{B}) \leq \Lambda(\mathcal{B}_0) \leq \lambda . \quad (5.3)$$

**Proposition 5.5**

Let  $\lambda = \lambda_k$  be an eigenvalue  $H(\Omega)$ . Then

$$\lambda_k \leq \inf_{\mathcal{B} \in \mathfrak{D}_k} \Lambda(\mathcal{B}). \quad (5.4)$$

**Proof of Proposition 5.5.**

The basic idea (which is already present in Courant's Theorem) is simply the following. We can assume (using Proposition 5.1)

$$\lambda_{k-1} < \lambda_k .$$

Attached to a minimal  $\mathcal{B}_k$  (hence regular), we have a  $k$ -dimensional space in  $H_0^1(\Omega)$  generated by the ground states of the  $D_i$  ( $i = 1, \dots, k$ ). We can find in this space a non trivial element which is orthogonal to the eigenspace corresponding to the eigenvalues which are  $\leq \lambda_{k-1}$ , whose energy is  $\mathfrak{L}_k$ , hence by the Minimax Principle  $\lambda_k \leq \mathfrak{L}_k$ .

Suppose now that we have the equality  $\lambda_k = \mathfrak{L}_k$ . Again by the proof of the Minimax Principle, this non trivial element should be an eigenfunction which is consequently Courant-sharp and we have consequently  $\lambda_k = L_k = \mathfrak{L}_k$ .

The following corollary is just a rephrasing of Propositions 5.5 and 5.4.

**Corollary 5.6**

We have

$$L_k \geq \mathfrak{L}_k \geq \lambda_k , \forall k \geq 1 . \quad (5.5)$$

In particular, if  $L_k = \lambda_k$  (also called the Courant-sharp case see in [4]) the nodal domain of a corresponding eigenfunction gives a minimal partition.

**Remarks 5.7**

(i) For the one dimensional case the standard Sturm-Liouville theory leads easily to the following

$$L_k = \mathfrak{L}_k = \lambda_k , \forall k \geq 1 . \quad (5.6)$$

(ii) It is easy to show, that for a given  $H$

$$\mathfrak{L}_1 = L_1 = \lambda_1, \tag{5.7}$$

(by the property of the ground state) and that

$$\mathfrak{L}_2 = L_2 = \lambda_2, \tag{5.8}$$

by the orthogonality of  $u_2$  to the ground state combined with Courant's nodal Theorem. (See also [9], Corollary 4.1 (Case  $V = 0$ )), but the extension to  $V \in L^\infty$  is not a problem.

(iii) The sequence  $L_k$  is not necessarily monotone : see for example (9.6).

One also observes that, using (5.5) and the property that  $\lambda_k \rightarrow +\infty$ ,

$$\lim_{k \rightarrow +\infty} \mathfrak{L}_k = +\infty . \tag{5.9}$$

## 6 Playing around Pleijel's argument.

It is a well known result of Pleijel that the one dimensional result that the  $k$ -th eigenvector of a Sturm-Liouville operator on an interval has only  $k$  nodal domains cannot be extended to higher dimension. The  $k$ -th Courant-sharp eigenfunctions (i.e.  $k$ -th eigenfunctions with  $k$  nodal domains) can only be found for a finite number of  $k$ 's.

We will show in this section, that the arguments behind the proof of this theorem give also many informations on the spectral minimal partition sequence in comparison with the spectral sequence and the nodal sequence.

Let us look at a universal lower bound for  $\mathfrak{L}_k(\Omega)$ . We actually obtain :

### Proposition 6.1

*Considering the Dirichlet Laplacian, we have*

$$\mathfrak{L}_k(\Omega) \geq k \frac{\pi j^2}{|\Omega|} ,$$

where  $|\Omega|$  denotes the area of  $\Omega$  and  $j$  is the smallest positive 0 of the Bessel function  $J_0$  :

$$j \sim 2.4048 \dots . \tag{6.1}$$

### Proof

For any  $D_j$  of a partition, we have by the Faber-Krahn inequality

$$|D_j| \lambda(D_j) \geq \pi j^2 .$$

The Faber-Krahn inequality gives indeed :

$$\lambda(D) \geq \frac{\lambda(B_{1/\sqrt{\pi}})}{|D|} , \tag{6.2}$$

for any open set  $D$ .

The lowest eigenvalue for the disk of radius 1 is known to be :

$$\lambda(B_1) = j^2 , \text{ with } \pi j^2 \sim 18.1695 . \tag{6.3}$$

Summing up over  $j$ , we obtain

$$\pi j^2 k \leq \sum_j |D_j| \lambda(D_j) \leq |\Omega| \max \lambda(D_j) . \tag{6.4}$$

Taking the infimum over the partition leads to the result.

**Remark 6.2**

Using Corollary 5.6, this implies

$$L_k(\Omega) \geq k \frac{\pi j^2}{|\Omega|} .$$

We conclude this section with a classical result of Pleijel, [23]:

**Theorem 6.3**

If  $\Omega$  is bounded and smooth,

$$\{k \mid u_k \text{ has } k \text{ nodal domains}\} \text{ is finite.} \quad (6.5)$$

This holds in larger generality for bounded potentials and also for higher dimensions.

Let us describe for completeness how Pleijel's Theorem is proved. The Weyl theory says that

$$\lambda_n \sim \frac{4\pi n}{|\Omega|} , \quad (6.6)$$

as  $n \rightarrow +\infty$ .

If  $u_n$  is an eigenfunction associated to  $\lambda_n$  with  $n$  nodal domains, we obtain immediately a contradiction for  $n$  large between (6.6) and (6.4) (applied with the family of nodal domains of  $u_n$ ), having in mind the value of  $j$  given in (6.1). So  $u_n$  cannot have  $n$  nodal domains ! More precisely, if there exists a smallest  $n(k)$  such that  $\lambda_{n(k)} = L_k$ , we obtain asymptotically

$$\liminf_{k \rightarrow +\infty} \frac{n(k)}{k} \geq j^2/4 > 1 . \quad (6.7)$$

A more difficult question is to determine whether  $L_k$  is always finite.

**Proposition 6.4**

In the case of the Laplacian and if  $\Omega$  is regular, we have<sup>2</sup>

$$\limsup_{k \rightarrow +\infty} \frac{\mathfrak{L}_k(\Omega)}{k} \leq \lambda_1(Hx_1)/|\Omega| , \quad (6.8)$$

where  $Hx_1$  is the regular hexagon of area 1.

---

<sup>2</sup>Thanks to M. Van den Berg for discussions. In particular he conjectures the existence of the limit  $\lim_{k \rightarrow +\infty} \frac{\mathfrak{L}_k}{k}$  and that the limit is actually  $\lambda_1(Hx_1)$ .

For the proof, we just use a (non strong) partition of  $\Omega$  by equal hexagons of area at most  $\frac{|\Omega|}{k}$ .

**Remark 6.5 .**

*Adding a potential does not create any difficulty and the previous discussions can be easilily adapted to go from  $H_0(\Omega)$  to  $H(\Omega)$ . Concerning the values  $L_k$  we obtain immediately,*

$$L_k^V(\Omega) \geq k \frac{\pi j^2}{|\Omega|} - \sup |V| . \quad (6.9)$$

*On the other hand, using the minimax, there are no problem to show that*

$$\mathfrak{L}_k^V(\Omega) \geq k \frac{\pi j^2}{|\Omega|} - \sup |V| . \quad (6.10)$$

*This implies in particular that*

$$\liminf_{k \rightarrow +\infty} \frac{\mathfrak{L}_k^V(\Omega)}{k} \geq \frac{\lambda_1(B_{1/\sqrt{\pi}})}{|\Omega|} > \frac{4\pi}{|\Omega|} , \quad (6.11)$$

*is satisfied in full generality.*

## 7 Minimal partitions and Courant-sharp

The main object of this section is the proof of Theorem 1.17.

### 7.1 Preliminaries

Except possibly the question of eliminating the assumption that  $\Omega$  is simply connected, the proof in this section uses essentially [16, 17] (or easy extensions of it). The minimal partitions which are involved in the proof are indeed regular. Although not very important here, this seems useful to mention for possible extensions in higher dimensions where we do not have the fine results established in Section 4.

### 7.2 Definition of an exhausting family $N(u, \alpha)$ .

Let  $u$  be an eigenfunction of  $H(\Omega)$  with  $k$  nodal domains and consider  $N(u) \in \mathcal{M}(\Omega)$ . First we consider the finite sets of points

$$C^*(N) := \mathcal{Z}_3 \cup (N(u) \cap \partial\Omega). \quad (7.1)$$

From each of these points an arc emanates which ends either in the point itself (loop) or ends in another point in  $C^*(N)$ . We call the collection of these arcs  $\mathcal{A}_*$ . Then we consider those components of  $N(u)$  whose intersection with  $C^*(N)$  is empty. They have to be pairwise disjoint imbedded circles (without selfintersections) and we call the collection of these circles  $\mathcal{A}_{**}$ . Let us introduce

$$\mathcal{A} = \mathcal{A}_* \cup \mathcal{A}_{**}.$$

Note that each arc (or loop)  $A \in \mathcal{A}$  is rectifiable (because  $N(u_m)$  is regular by Theorem 2.3) and we can associate to  $A$  naturally a middle point  $x_A$  in the natural way ( $x_A$  is chosen arbitrarily if  $A \in \mathcal{A}_{**}$ ). We have a natural arc length parametrization starting from the point  $x_A$ , but we prefer to parametrize  $A$  as a parametrized curve  $[-1, +1] \ni t \mapsto L(A, t)$  such that

$$L(A, 0) = x_A, \quad L(A, -1) = y_A^-, \quad L(A, 1) = y_A^+,$$

where  $y_{A^-}$  and  $y_{A^+}$  are the end points in the case of an arc, and where  $y_{A^-} = y_{A^+}$  is the irregular point in the case of an irregular loop (i.e. in  $\mathcal{A}_*$ ) and the opposite point in the case of a regular loop (i.e. in  $\mathcal{A}_{**}$ ).

For each  $\alpha \in (0, 1)$ , we can consider the set

$$N(u, \alpha) = \{N(u) \setminus L(A, (-1 + \alpha, 1 - \alpha))\} \quad (7.2)$$

and complete the definition by

$$N(u, 0) = \emptyset \text{ and } N(u, 1) = N(u) . \quad (7.3)$$

Note that by construction for every  $0 < \alpha$ ,  $N(u, \alpha)$  contains all the critical points and  $N(u) \cap \partial\Omega$ ; this will be important below.

### 7.3 Proof of Theorem 1.17.

**We assume for contradiction that for some  $k$ ,  $\mathfrak{L}_k = L_k$ , but that  $\lambda_k < \lambda_m = L_k$  for some  $m > k$ .**

Taking the smallest  $m$  with this property, we can in addition assume that

$$\lambda_{m-1} < \lambda_m . \quad (7.4)$$

Let  $u_m$  a normalized eigenfunction such that  $u_m$  has  $\mu(u_m) = k$  nodal domains  $D_i$  ( $i = 1, 2, \dots, k$ ). Then we associate with the exhausting family  $N(u_m, \alpha)$  the decreasing family of open sets :

$$\Omega(\alpha) = \Omega \setminus N(u_m, \alpha) . \quad (7.5)$$

We want to consider the spectrum of

$$H(\alpha) := H(\Omega(\alpha)) . \quad (7.6)$$

(We suppress the dependence on  $\Omega$  and  $V$ .)

Then,  $H(0)$  is our initial operator  $H(\Omega)$  and

$$H(1) = \bigoplus_{i=1}^k H(D_i) \quad (7.7)$$

Hence  $H(1)$  has as lowest eigenvalue  $\lambda_m$  with multiplicity  $k$ . By construction  $\sigma(H(0)) = \sigma(H)$ . Furthermore  $\lambda_1(H(1))$  has multiplicity  $k$  and

$$\lambda_{k+1}(H(1)) > \lambda_k(H(1)) = \lambda_m(H(0)) .$$

#### Lemma 7.1

*For any  $\ell$ ,  $\lambda_\ell(H(\alpha))$  is monotonically increasing with  $\alpha$ .*

**Proof**

We just note that the form domains  $\mathcal{Q}(\alpha)$  of the quadratic forms  $q(\alpha)$  associated to  $H(\alpha)$ ,  $0 \leq \alpha \leq 1$ , satisfy  $\mathcal{Q}(\alpha) \subset \mathcal{Q}(\alpha')$  for  $\alpha' \leq \alpha$ .

**Lemma 7.2**

For any  $\ell$ ,  $\lambda_\ell(H(\alpha))$  depends continuously upon  $\alpha$ .

**Proof**

Although there is a lot of literature<sup>3</sup> on the subject (see [22, 26, 27, 30, 31, 32, 28]), it is difficult to give a reference corresponding to this crack situation. This statement is proved in Dauge-Helffer [10] (at least for the case of one crack). These authors treat the case when the condition on the crack is Neumann. The Dirichlet case does not create new problems (the important point being the monotonicity which is evident in the case of Dirichlet). Note that we have strong convergence and that the left continuity and the right continuity should be treated separately.

We continue with

**Lemma 7.3**

For each  $\alpha \in [0, 1]$ ,  $\lambda_m \in \sigma(H(\alpha))$ .

**Proof**

By construction of  $N(u_m, \alpha)$ , the restriction of  $u_m$  to  $\Omega(\alpha)$  is indeed an eigenfunction of  $H(\alpha)$ .  $\square$

We recall (see the two first lines of this subsection) that we are inside a proof by contradiction and consider first the following case.

**Case (a):  $\lambda_m$  is simple****Lemma 7.4**

There is a minimal  $\alpha_1 \in (0, 1)$  such that

$$\lambda_1(\alpha_1) < \lambda_2(\alpha_1) \leq \dots \leq \lambda_{m-1}(\alpha_1) = \lambda_m(\alpha_1), \quad (7.8)$$

with  $\lambda_m(\alpha_1) = \lambda_m$ .

---

<sup>3</sup>We thank P. Stollmann and M. Dauge for useful discussions.

**Proof**

By assumption  $m > k$  and  $H(0)$  has  $m$  eigenvalues smaller or equal to  $\lambda_m$  whereas  $H(1)$  has just  $k$  eigenvalues smaller or equal to  $\lambda_m$ , hence less. By continuity with respect to  $\alpha$  at least one eigenvalue has to become larger than  $\lambda_m$ .  $\square$

Next we consider this eigenvalue of  $H(\alpha_1)$ . The restriction of  $u_m$  to  $\Omega(\alpha_1)$  gives a first eigenfunction and there exists a real valued normalized eigenfunction  $v$  of  $H(\alpha_1)$  such that  $v$  is orthogonal to  $u_m$  in  $L^2(\Omega(\alpha_1))$  :

$$\langle u_m | v \rangle = 0 , \quad (7.9)$$

where  $\langle \cdot | \cdot \rangle$  denotes the  $L^2$ -scalar product.

We now play inside the two dimensional eigenspace spanned by  $u_m$  and  $v$ .

**Lemma 7.5**

*There exists  $\beta_0 > 0$ , such that  $\forall \beta \in (-\beta_0, +\beta_0)$  the function  $w_\beta = u_m + \beta v$  has exactly  $k$  nodal domains.*

*Furthermore, the family of the nodal domains of  $w_\beta$  gives, for  $\beta \neq 0$ , a minimal bipartite partition of  $\Omega$ , which is distinct of the partition associated to  $N(u_m)$ .*

**Proof**

We recall that this construction is done for  $\alpha = \alpha_1 \in (0, 1)$ . For each  $A \in \mathcal{A}$ , let

$$I_A := L(A, (-1 + \alpha_1, 1 - \alpha_1)) ,$$

and let  $\mathcal{V}_A \subset \Omega$  be an open neighborhood of  $\overline{I_A}$ , whose regular boundary crosses  $N(u_m)$  twice (transversally) and such that each component of the open set  $\Omega(\alpha_1) \setminus \cup_{A \in \mathcal{A}} \overline{\mathcal{V}_A}$  (which is contained in  $\Omega(1)$ ), is contained in a unique nodal domain of  $u_m$ .

We assume that we have colored these nodal domains by  $+$  or  $-$ , and this permits to write, for each  $A \in \mathcal{A}$ , the decomposition

$$\partial \mathcal{V}_A \cap \Omega(\alpha_1) = b_A^+ \cup b_A^- ,$$

where  $b_A^\pm$  is contained in a positive or negative nodal domain of  $u_m$ .

The first claim is now that there exists  $\beta_0$  such that if we add  $\beta v$ , with  $|\beta| \leq \beta_0$ , the number of nodal domains of  $u_m + \beta v$  can only increase. Using Hopf's boundary point lemma for  $u = u_m$ , [12], we have  $|\nabla u_m(x)| > 0$

for  $x \in (N(u_m) \setminus N(u_m, \alpha_1)) \cap \Omega$  and using the property that  $v$  vanishes at the boundary of  $\Omega(\alpha)$ , we obtain the existence of  $\beta_0$  such that  $u_m + \beta v$  is strictly positive on each  $b_A^+$  and strictly negative on  $b_A^-$ . It is then clear that associated to each positive  $D_i$ , there is at least one nodal domain of  $u_m + \beta v$ , with non trivial intersection with  $D_i$  and contained in  $D_i \cup (\cup_A \mathcal{V}_A)$ . All these nodal domains are necessarily disjoint and this proves the first claim.

Let us now show that we cannot increase the number of nodal domains. If it was the case, this would give an upper bound for  $\mathfrak{L}_{k+1}$  and using (5.1), we would obtain, using the strict monotonicity of the sequence  $\mathfrak{L}_\ell$  (see (5.1)) with respect to  $\ell$ ,  $\lambda_m = \mathfrak{L}_k < \mathfrak{L}_{k+1} \leq \lambda_m$ , hence a contradiction.

So  $w_\beta$  has also exactly  $k$  nodal domains corresponding also to a minimal  $k$ -partition  $\mathcal{D}' \in \mathfrak{D}_k$  of  $\Omega$ . But  $\mathcal{D}' \neq \mathcal{D}$  since both functions,  $u_m$  and  $v$  (and hence  $w_\beta$  for  $\beta \neq 0$ ) are linearly independent. In addition, one can verify that  $G(\mathcal{D}')$  is bipartite.  $\square$

Now we can complete the proof of Theorem 1.17 for **case (a)**. Indeed by Theorem 1.14,  $w_\beta$  (more precisely the natural extension of  $w_\beta$  to  $\Omega$ ) is an eigenfunction of  $H = H(0)$  and therefore this would imply that  $\lambda_m$  has multiplicity at least two, contradicting our assumption that  $\lambda_m$  is simple.

### Remark 7.6

*Note that for these proofs we only need weak versions of our results because we work only with strong regular partitions (satisfying in addition the equal angle meeting property). So the techniques of [16] (as recalled in appendix B) are also relevant.*

*In the non simply connected case, if one wants to apply [16], one should also verify a global compatibility condition for each homotopy class of  $\Omega$ . Because  $w_\beta$  is an eigenfunction in  $\Omega(\alpha_1)$ , this is a consequence of the property that any path in  $\Omega$  is homotopic to a path in  $\Omega(\alpha_1)$ . It is then easy to verify this additional cycle-compatibility condition introduced in [16], because  $w_\beta$  is an eigenfunction of  $H(\alpha_1)$ .*

### Case (b): $\lambda_m$ has multiplicity greater than one.

Assume that  $\lambda_m$  has multiplicity  $\ell > 1$ , so that the  $m + \ell$  first eigenvalues of  $H = H(0)$  satisfy :

$$\lambda_1 < \lambda_2 \leq \dots \leq \lambda_k \leq \dots \lambda_{m-1} < \lambda_m = \dots = \lambda_{m+\ell-1} < \lambda_{m+\ell} . \quad (7.10)$$

Of course, all the previous constructions can be done but arriving at the last line of case (a), we loose the contradiction. The idea is that we have to choose our  $w$  more carefully.

$\lambda_m$  being an eigenvalue of  $H(\alpha)$  for any  $\alpha$ , we can associate to  $\lambda_m$  the eigenspace  $\underline{U}(\alpha, \lambda_m)$  which is defined as the subspace in the spectral space of  $H(\alpha)$   $U(\alpha, \lambda_m)$  consisting of functions which are restrictions to  $\Omega(\alpha)$  of eigenfunctions of  $H(\Omega)$ . Of course  $\underline{U}(\alpha, \lambda_m)$  contains  $u_m$  but could be larger. We then need to show that

**Lemma 7.7**

For  $\alpha = \alpha_1$ ,  $\underline{U}(\alpha_1, \lambda_m)$  is strictly included in  $U(\alpha_1, \lambda_m)$ .

**Proof**

For any  $\alpha < \alpha_1$ , we can choose a normalized  $v_\alpha$  in  $U(\alpha, \lambda_{m-1}(\alpha))$ . Then it is clear, observing that  $\lambda_{m-1}(\alpha) < \lambda_m$ , that

- $v_\alpha$  is orthogonal in  $L^2(\Omega(\alpha))$  to  $\underline{U}(\alpha_1, \lambda_m)$  which is (more precisely, can be identified to) a subspace of  $U(\alpha, \lambda_m)$  for any  $\alpha < \alpha_1$ .
- $v_\alpha$  is bounded independently of  $\alpha$  in  $H^1(\Omega(\alpha_1))$ .

Then we can by compactness, find a sequence  $w_{\alpha(n)}$  such that  $\alpha(n)$  tends to  $\alpha_1$  as  $n \rightarrow +\infty$  and  $v_{\alpha(n)}$  converges weakly to some  $v_{\alpha_1}$  in  $W^1(\Omega(\alpha_1))$  and strongly in  $W^s(\Omega(\alpha_1))$  for  $s < 1$  by compactness.

Now, it is clear that

- $v_{\alpha_1}$  is orthogonal to  $\underline{U}(\alpha_1, \lambda_m)$ .
- $\|v_{\alpha_1}\| = 1$ .
- $(-\Delta + V)v_{\alpha_1} = \lambda_m v_{\alpha_1}$  in  $\Omega(\alpha_1)$ .

With a small additional work, one can show that  $v_{\alpha_1} \in W_0^1(\Omega(\alpha_1))$ . So  $v_{\alpha_1}$  is effectively in the form domain for the Dirichlet problem in  $\Omega(\alpha_1)$  and in  $U(\alpha_1, \lambda_m) \cap \underline{U}(\alpha_1, \lambda_m)^\perp$ .

**End of the proof of case (b).**

The argument is then as in case (a), but, using Lemma 7.7, we can choose a non trivial  $v$  in  $U(\alpha_1, \lambda_m) \setminus \underline{U}(\alpha_1, \lambda_m)$ . But on one hand  $w_\beta$  cannot belong to  $\underline{U}(\alpha_1, \lambda_m)$  (because  $v$  does'nt). On the other hand, we have obtained some  $\beta \neq 0$  such that  $w_\beta$  extends as an eigenfunction of  $H(\Omega)$  hence by definition in  $\underline{U}(\alpha_1, \lambda_m)$  and this gives the contradiction.

Theorem 1.17 has an immediate consequence.

**Corollary 7.8**

*Let  $\Omega$  satisfy Assumption 1.1 and assume that  $V \in L^\infty(\Omega)$ . Then*

$$|\{k \mid \mathfrak{L}_k = L_k\}| < \infty. \tag{7.11}$$

**Proof**

This is an immediate consequence of Theorem 1.17 and of Pleijel's Theorem 6.3. □

**Remark 7.9**

*It is now easier to analyze the situation for the disk and for rectangles (at least in the irrational case), since we have just to check for which eigenvalues one can find associated Courant-sharp eigenfunctions. This will be done in sections 9 and 10.*

## 8 Further properties of subpartitions

All the statements of this section illustrate the rigidity of the structure of the subpartitions. This can be very efficient for disproving that a partition is minimal. In particular we will prove Theorem 1.13.

The following proposition is useful :

### Proposition 8.1

Under Assumptions 1.1 and 1.2, let  $\mathcal{D} = (D_i)_{i=(1,\dots,k)}$  be a  $k$ -minimal partition for  $\mathfrak{L}_k(\Omega)$ . Then, for any subset  $I \in \{1, \dots, k\}$ , the associated subpartition  $\mathcal{D}^I = (D_i)_{i \in I}$  satisfies

$$\mathfrak{L}_k = \Lambda(\mathcal{D}^I) = \mathfrak{L}_{|I|}(\Omega^I), \quad (8.1)$$

where

$$\Omega^I := \text{Int}(\overline{\cup_{i \in I} D_i}).$$

### Proof

We prove this proposition by contradiction. If it was not the case, we would construct (starting of a minimal  $|I|$ -partition of  $\Omega^I$ ) a new minimal partition  $\tilde{\mathcal{D}}$  of  $\Omega$ , for which the  $\lambda(\tilde{D}_i)$ 's are not equal in contradiction with what we proved in Section 4 (see also (d) in Remark 3.7 together with the fact that  $k_0 = \emptyset$  in our case).

As a consequence of a more general theorem in [4], we have (with very weak assumptions on  $\Omega$ ) the analogous of Proposition 8.1

### Proposition 8.2

Suppose  $u$  is **Courant-sharp**. Denote the associated nodal domains by  $\{D_i\}_1^k$ . Let  $L$  be a subset of  $\{1, 2, \dots, k\}$  with  $\#L = \ell < k$  and let  $\Omega^L = \text{Int}(\overline{\cup_{i \in L} D_i}) \setminus \partial\Omega$ . Then

$$\lambda_\ell(\Omega^L) = \lambda_k \quad (8.2)$$

where  $\lambda_j(\Omega^L)$  are the eigenvalues of  $H(\Omega^L)$ .

Moreover, if  $\Omega^L$  is connected,  $u|_{\Omega^L}$  is **Courant-sharp** and  $\lambda_\ell(\Omega^L)$  is simple.

### Proposition 8.3

Under the assumptions of the introduction on  $\Omega$  and  $V$ , let  $\mathcal{D} = (D_1, \dots, D_k)$

be a minimizing partition relative to  $\mathfrak{L}_k$ . Let  $\mathcal{D}' \subset \mathcal{D}$  be any subpartition into  $1 \leq k' \leq k$  elements which is bipartite relatively to

$$\Omega' := \text{Int} \left( \bigcup_{D_i \in \mathcal{D}'} \overline{D_i} \right).$$

Then

- (a)  $\mathfrak{L}_k = \lambda_{k'}(\Omega')$ ;
- (b) If  $k' < k$ , and  $\Omega'$  is connected then  $\lambda_{k'}(\Omega')$  is simple.

**Proof**

The point (a) is proven like at the end of Section 4. The proof of (b) is an immediate consequence of Proposition 8.2, if we add the assumption that there exists a bipartite subpartition  $\mathcal{D}''$  such that  $\mathcal{D}' \subset \mathcal{D}'' \subset \mathcal{D}$  with  $k' < k'' \leq k$ .

The general proof is a little more tricky. Given the minimal partition  $\mathcal{D}$  of  $\Omega$ , and the subpartition  $\mathcal{D}'$  (with associated bipartite graph), there is a subfamily  $\mathcal{A}_0$  of the set  $\mathcal{A}$  of arcs (we keep the ends of the arcs living in  $\Omega$ ) belonging to  $N(\mathcal{D})$  (see the discussion in Section 7) such that  $\tilde{\Omega} := \Omega \setminus \mathcal{A}_0$  has for the same partition a bipartite graph relatively to  $\tilde{\Omega}$  and such that  $\mathcal{A}_0$  does not contain arcs belonging to the intersection of the boundaries of two neighbors of  $\mathcal{D}'$  in  $\Omega'$ . But we are in a Courant-sharp situation, so

$$\mathfrak{L}_k(\Omega) = \mathfrak{L}_k(\tilde{\Omega}) = \lambda_k(\tilde{\Omega}), \tag{8.3}$$

and we can apply the Courant-sharp Theorem relative to  $\tilde{\Omega}$  and the partition  $\mathcal{D}$  (which is now the family of nodal domains of an eigenfunction on  $\tilde{\Omega}$ ).

**Remark 8.4**

*In the preceding proof,  $\tilde{\Omega}$  is not unique, but it is interesting to emphasize that what we have shown is that, once a minimal  $k$ -partition of  $\Omega$  is given, then all the possible  $\tilde{\Omega}$ 's should share the property (8.3).*

In order to complete the proof of Theorem 1.13, it remains to establish our uniqueness result at the level of the subpartitions of a minimal partition.

**Proposition 8.5 (Uniqueness)**

*Let  $\mathcal{D}$  be a minimal  $k$ -partition relative to  $\mathfrak{L}_k(\Omega)$ . Let  $\mathcal{D}' \subset \mathcal{D}$  be any subpar-*

tion of  $\mathcal{D}$  into  $1 \leq k' < k$  elements and let

$$\Omega' = \text{Int} \left( \bigcup_{D_i \in \mathcal{D}'} \overline{D_i} \right),$$

be connected. Then  $\mathfrak{L}_{k'}(\Omega')$  is uniquely achieved.

We know already from Proposition 8.1 that  $\mathfrak{L}_{k'}(\Omega') = \mathfrak{L}_k(\Omega)$ . The proof of uniqueness is by contradiction. Let  $I$  and  $J$  two subsets of  $\{1, \dots, k\}$  such that  $I \cap J = \emptyset$  and  $I \cup J = \{1, \dots, k\}$ .

If we have indeed two minimal subpartitions of  $\Omega_I$  for some  $I$  of cardinality strictly less than  $k$ . We can complete these two subpartitions by the open sets  $D_j$  ( $j \in J$ ). According to the proof of our results. Now take a pair  $(D_i, D_j)$ , with  $i \sim j$  ( $i \in I$  and  $j \in J$ ). This should exist if  $\Omega$  is connected and  $\Omega_I$  is connected. There is necessarily another  $\tilde{D}_i$  of the other partition which meets  $D_i$  and is a neighborhood of  $D_j$ . But the eigenfunction  $u_{ij}$  of  $D_{ij}$  and  $\tilde{u}_{ij}$  of  $\tilde{D}_{ij}$  should be proportional on  $D_j$  to  $u_j$ . It is then clear that by unique continuation  $D_i$  should coincide with  $\tilde{D}_i$  and  $u_i$  should be proportional to  $\tilde{u}_i$ . Possibly iterating the argument, we arrive to a contradiction that the two partitions are different.

## 9 Example 1: The case of the disk

In this section, we analyze in great detail the case of the disk. Although the spectrum is explicitly computable, we are mainly interested in the ordering of the eigenvalues corresponding to different angular momenta. In particular this will give a first example where the inequalities in (5.5) can be strict.

Consider the Dirichlet realization  $H_0$  in the unit disk  $B_1 \subset \mathbb{R}^2$ . We have in polar coordinates :

$$-\Delta = -\frac{\partial^2}{\partial r^2} - \frac{1}{r} \frac{\partial}{\partial r} - \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} ,$$

and the Dirichlet boundary conditions require that any eigenfunction  $u$  satisfies  $u(r, \theta) = 0$  for  $r = 1$ . We analyze for any  $\ell \in \mathbb{N}$  the eigenvalues  $\lambda_{\ell, j}$  of

$$\left(-\frac{d^2}{dr^2} - \frac{1}{r} \frac{d}{dr} + \frac{\ell^2}{r^2}\right) f_{\ell, j} = \lambda_{\ell, j} f_{\ell, j} , \quad \text{in } (0, 1) .$$

We observe that the operator is self adjoint for the scalar product in  $L^2((0, 1), r dr)$ . The corresponding eigenfunctions of the eigenvalue problem take the form

$$u(r, \theta) = f_{\ell, j}(r) (a \cos \ell \theta + b \sin \ell \theta) , \quad \text{with } a^2 + b^2 > 0 , \quad (9.1)$$

where the  $f_{\ell, j}(r)$  are suitable Bessel functions satisfying for  $\ell = 0$ ,  $f'_{0, j}(0) = 0$  and  $f_{0, j}(1) = 0$  and for  $\ell > 0$ ,  $f_{\ell, j}(0) = f_{\ell, j}(1) = 0$ . For the corresponding  $\lambda_{\ell, j}$ 's, we find by looking up for instance [5] (or in appendix A) the following ordering.

$$\begin{aligned} \lambda_1 = \lambda_{0,1} &< \lambda_2 = \lambda_3 = \lambda_{1,1} < \lambda_4 = \lambda_5 = \lambda_{2,1} < \lambda_6 = \lambda_{0,2} < \lambda_7 = \lambda_8 = \lambda_{3,1} < \dots \\ &\dots < \lambda_9 = \lambda_{10} = \lambda_{1,2} < \lambda_{11} = \lambda_{12} = \lambda_{4,1} < \dots \\ &\dots < \lambda_{13} = \lambda_{14} = \lambda_{2,2} < \lambda_{15} = \lambda_{0,3} < \dots \end{aligned} \quad (9.2)$$

We recall that the zeros of the Bessel functions are related to the eigenvalues by the relation

$$\lambda_{\ell, k} = (j_{\ell, k})^2 . \quad (9.3)$$

We hence have from (9.1)

$$\begin{aligned}
\mu(u_1) &= 1, \\
\mu(u) &= 2, \text{ for any eigenfunction } u \text{ associated to } \lambda_2 = \lambda_3, \\
\mu(u) &= 4, \text{ for any eigenfunction } u \text{ associated to } \lambda_4 = \lambda_5, \\
\mu(u_6) &= 2, \\
\mu(u) &= 6, \text{ for any eigenfunction } u \text{ associated to } \lambda_7 = \lambda_8, \\
\mu(u) &= 4, \text{ for any eigenfunction } u \text{ associated to } \lambda_9 = \lambda_{10}, \\
\mu(u) &= 8, \text{ for any eigenfunction } u \text{ associated to } \lambda_{11} = \lambda_{12}, \\
\mu(u) &= 8, \text{ for any eigenfunction } u \text{ associated to } \lambda_{13} = \lambda_{14}, \\
\mu(u_{15}) &= 3.
\end{aligned} \tag{9.4}$$

Hence

$$L_1 = \lambda_1, L_2 = \lambda_2, L_3 = \lambda_{15}, L_4 = \lambda_4 \tag{9.5}$$

and this implies

$$L_3 > \lambda_6 > L_4 > \lambda_3. \tag{9.6}$$

In addition, let us show that

$$L_3 > \mathfrak{L}_3. \tag{9.7}$$

This can be seen as follows. We can split  $B_1$  in three sectors with opening angle  $2\pi/3$ . Call such a sector  $S_{1/3}$  then the corresponding eigenvalue  $\lambda(S_{1/3})$  satisfies :

$$\mathfrak{L}_3 \leq \lambda(S_{1/3})$$

We then observe that by monotonicity that :

$$\lambda(S_{1/3}) < \lambda(S_{1/4})$$

and we can recognize that  $\lambda(S_{1/4}) = \lambda_4$  observing that  $\lambda_4 = \lambda_{\ell,j}$  with  $\ell = 2$  and  $j = 1$  (see (9.1)). The proof of (9.7) is then a consequence of (9.6).

**Remark 9.1**

*An open problem is to prove or disprove the equality  $\mathfrak{L}_3 = \lambda(S_{1/3})$ . A possibility for trying to find other configurations could be to look at the second eigenvalue of a half disk problem  $\{(x_1^2 + x_2^2) < 1\} \cap \{x_2 > 0\}$ , where we take Dirichlet on the circular part and a part of the straight basis and Neumann on the other part. More precisely, we take Neumann on  $x_2 = 0$ ,  $|x_1| < t$ , where  $t$  is a free parameter. The nodal domains of the second eigenfunction (completed by symmetry) could give an alternative candidate for such a minimal partition.*

In the case of the disk, we have

**Proposition 9.2**

*Except the cases  $k = 1, 2$  and  $4$ , minimal partitions never correspond to nodal domains.*

**Proof**

According to Theorem 1.17, it is enough to investigate when the  $k$ -th eigenvalue corresponds to Courant-sharp eigenfunctions.

One can in addition use the twisting trick (see [18]) for eliminating all the eigenvalues  $\lambda_{\ell,m}$ , for which  $m \geq 2$  and  $\ell > 0$ . This trick goes roughly as follows. When  $\ell > 0$ , we can divide the disk as the union of a smaller disk and of its complementary, each of these sets being the union of at least two nodal domains. Then by small rotation of the small disk, we obtain a new partition which has the same energy. If the initial one was minimal, the new one should be also minimal, but it is easy to show that the new one has not the “equal angle meeting” property of a regular partition. This gives the contradiction.

So we have finally to analyze the eigenvalues  $\lambda_{0,k}$  and the family  $\lambda_{\ell,1}$ .

For the first family, we observe that  $\lambda_{0,k}$  can neither be the  $k$ -th eigenvalue as soon as  $k \geq 2$ .

For the second family, which occurs only for  $k = 2\ell$  even, inspection of the tables leads to the condition  $k \leq 4$ , we observe indeed that  $\lambda_{0,2} < \lambda_{3,1}$ .

We also observe that when  $k$  is odd, we obtain that necessarily  $L_k = \lambda_{0,k}$ .

**Remark 9.3**

*It could be interesting to determine when  $\mathfrak{L}_k$  is an eigenvalue of the Laplacian on the double covering. Again, one can show (see [18]) that this cannot be the case for  $k$  large.*

## 10 Example 2: the case of the rectangle

Note that for the case of a rectangle, the spectrum and the properties of the eigenfunctions are analyzed as toy models in [23], Section 4. This was also used for testing general conjectures in [4].

For a rectangle of sizes  $a$  and  $b$ , the spectrum is given by  $\pi^2(m^2/a^2 + n^2/b^2)$   $((m, n) \in (\mathbb{N}^*)^2)$ .

The first remark is that all the eigenvalues are simple if  $\frac{a^2}{b^2}$  is irrational. Except for specific remarks for the square, we now assume

$$(a/b)^2 \text{ is irrational.}$$

So we can associate to each eigenvalue  $\lambda_{m,n}$ , an (essentially) unique eigenfunction  $u_{m,n}$  such that  $\mu(u_{m,n}) = nm$ . Given  $k \in \mathbb{N}^*$ , the lowest eigenvalue corresponding to  $k$  nodal domains is given by

$$L_k = \pi^2 \inf_{mn=k} (m^2/a^2 + n^2/b^2) .$$

The behavior of  $L_k$  can depend dramatically of the arithmetical properties of  $k$  but what is important for us is that in any case we have

$$L_k \geq 2\pi^2 k / (ab) . \tag{10.1}$$

This immediately implies

$$\liminf_{k \rightarrow +\infty} \frac{L_k}{k} \geq \frac{2\pi^2}{ab} . \tag{10.2}$$

We note that the right hand side can also be written in the form  $\lambda([0, 1]^2)/|\Omega|$ .

### Remark 10.1

*Note that these estimates are much better than the estimates obtained by Faber-Krahn inequality (see in Section 6).*

As we have seen in (6.8) and using<sup>4</sup> the comparison between the lowest eigenvalue of the hexagon and of the square,

$$\lambda(Hx1) \sim 18.59013 < \lambda([0, 1]^2) = 2\pi^2 \sim 19.7392 , \tag{10.3}$$

---

<sup>4</sup>We thank V. Bonnaillie-Noël and G. Vial for giving us a precise numerical approximation of  $\lambda(Hx1)$ . According to A. El Soufi, it seems unknown that the hexagon gives the minimal eigenvalue between all the polygons (of same area) permitting to realize a perfect partition of the plane. We just compare here the square and the hexagon.

this will imply

$$\liminf_{k \rightarrow +\infty} \frac{L_k}{k} > \limsup_{k \rightarrow +\infty} \frac{\mathfrak{L}_k}{k}. \quad (10.4)$$

This implies that  $L_k > \mathfrak{L}_k$  for  $k$  large.

**Remark 10.2**

*In the case when  $(\frac{a}{b})^2$  is rational we could have problems in the case of multiplicities. We have then to control the nodal sets of the eigenfunctions corresponding to the degenerate eigenvalues which are  $\leq \mathfrak{L}_k$ .*

We now describe all the possible situations.

**Lemma 10.3**

*In the irrational case,  $\lambda_{m,n}$  cannot lead to a Courant-sharp situation if  $\inf(m, n) \geq 3$ .*

**Proof**

Applying Proposition 8.2, it is sufficient to analyze the case when  $m = n = 3$ . It is then enough to show that  $\lambda_{3,3}$  cannot be the ninth eigenvalue. Because the eigenvalues corresponding to  $\max(m, n) \leq 3$  are obviously below, let us assume by contradiction that  $\lambda_{3,3} \leq \lambda_{1,4}$  and that  $\lambda_{3,3} \leq \lambda_{4,1}$ .

This reads

$$\frac{9}{a^2} + \frac{9}{b^2} \leq \frac{1}{a^2} + \frac{16}{b^2},$$

and

$$\frac{9}{a^2} + \frac{9}{b^2} \leq \frac{1}{a^2} + \frac{16}{b^2}.$$

So, we obtain

$$\frac{8}{7} \leq \frac{a^2}{b^2} \leq \frac{7}{8},$$

Hence a contradiction.

The next step is given in

**Lemma 10.4**

*In the irrational case,  $\lambda_{m,n}$  cannot lead to a Courant-sharp situation if  $m = 2$  and  $n \geq 4$  or if  $m \geq 4$  and  $n = 2$ .*

Again it is enough to look at the case  $(m = 2, n = 4)$ , and to show that it cannot be the eight-th eigenvalue. Using the same idea as in the previous lemma, we assume by contradiction that  $\lambda_{2,4} \leq \lambda_{1,5}$  and that  $\lambda_{2,4} \leq \lambda_{3,1}$ .

This reads

$$\frac{4}{a^2} + \frac{16}{b^2} \leq \frac{1}{a^2} + \frac{25}{b^2},$$

and

$$\frac{4}{a^2} + \frac{16}{b^2} \leq \frac{9}{a^2} + \frac{1}{b^2}.$$

So, we obtain

$$\frac{1}{3} \leq \frac{a^2}{b^2} \leq \frac{1}{3}.$$

But this gives  $\frac{a^2}{b^2} = \frac{1}{3}$ , which is excluded by the assumption that  $\frac{a^2}{b^2}$  is irrational.

Let us now analyze the remaining cases.

**In the case  $m = 2, n = 3$ ,** an eigenfunction corresponding to  $\lambda_{m,n}$  is Courant-sharp if

$$\frac{4}{a^2} + \frac{9}{b^2} \leq \frac{9}{a^2} + \frac{1}{b^2},$$

and

$$\frac{4}{a^2} + \frac{9}{b^2} \leq \frac{1}{a^2} + \frac{16}{b^2}.$$

So, we obtain

$$\frac{8}{5} \leq \frac{a^2}{b^2} \leq \frac{5}{3}.$$

**The case  $m = 3, n = 2$ ,** is obtained by exchanging the role of  $a$  and  $b$ . So, we obtain

$$\frac{8}{5} \leq \frac{b^2}{a^2} \leq \frac{5}{3}.$$

**In the case  $m = 2, n = 2$ ,** we obtain similarly

$$\frac{3}{5} \leq \frac{a^2}{b^2} \leq \frac{5}{3}.$$

**For the case  $m = 1, n = k$ ,** we obtain

$$\frac{1}{a^2} + \frac{k^2}{b^2} \leq \frac{4}{a^2} + \frac{1}{b^2},$$

So, we obtain simply


$$\frac{(k^2 - 1)}{3} < \frac{a^2}{b^2}.$$

Finally, **the case**  $m = k$ ,  $n = 1$  leads to the condition

$$\frac{(k^2 - 1)}{3} < \frac{b^2}{a^2}.$$

### A candidate for the 3-minimal partition on the square.

In the case of the square, an argument similar to the case of the disk shows that  $\mathfrak{L}_3$  (which should be smaller than  $\mathfrak{L}_4$ ) is strictly less than  $L_3$ . We observe indeed that  $\lambda_4$  is Courant-sharp, so  $\mathfrak{L}_4 = \lambda_4$ , and there is no eigenfunction corresponding to  $\lambda_2 = \lambda_3$  with three nodal domains (by Courant's Theorem).



Restricting to the half-rectangle and assuming that there is a minimal partition which is symmetric with one of the symmetry axes of the square perpendicular to two opposite sides, one is reduced to analyze a family of Dirichlet-Neumann problems. Numerical computations<sup>5</sup> performed by V. Bonnaillie-Noel (in January 2006) and G. Vial lead to a natural candidate (See Picture above) for a symmetric minimal partition.

The complete structure is reobtained from the half square by symmetry with respect to the horizontal axis. We observe numerically that the three lines of  $N(\mathcal{D})$  meet at the center of the square. As expected by the theory they meet with equal angle  $\frac{2\pi}{3}$  and start from the boundary orthogonally.

---

<sup>5</sup>See <http://www.bretagne.ens-cachan.fr/math/people/virginie.bonnaillie/BH/MinimalPartitions/>

# A Zeros of Bessel functions

Let  $j_{\ell,k}$  the  $k$ -th zero of the Bessel function corresponding to the integer  $\ell \in \mathbb{N}$ . The reference is the book by G.N. Watson [29]. The most important statement for us is that  $j_{\ell,k} = j_{\ell',k'}$  imply if  $\ell$  and  $\ell'$  are positive integers, that  $\ell = \ell'$  and  $k = k'$ . We refer to the subsection 15.28 (p. 484-485). Note that the proof of this result is based on deep results by Siegel about algebraic numbers.

Here is a list of approximate values after the celebrated handbook of [1], p. 409, we keep only the values which are less than approximately 13.

$\ell =$	0	1	2	3	4	5	6	7	8
$k = 1$	2.40	3.83	5.14	6.38	7.59	8.77	9.93	11.08	12.22
2	5.52	7.02	8.42	9.76	11.06	12.34	.		
3	8,65	10.17	11.62	13.02	..				
4	11.79	13.32	.						

(A.1)

This leads to the following ordering of the zeros :

$$\begin{aligned}
 j_{0,1} < j_{1,1} < j_{2,1} < j_{0,2} < j_{3,1} < j_{1,2} < j_{4,1} < j_{2,2} < j_{0,3} < \dots \\
 \dots < j_{5,1} < j_{3,2} < j_{6,1} < j_{1,3} < j_{7,1} < j_{2,3} < j_{0,4} < j_{8,1} .
 \end{aligned}
 \tag{A.2}$$

Note that, using Sturm-Liouville theory, (see for example the proof in [29]), the following inequalities are always true :

$$j_{\ell,k} < j_{\ell+1,k} < j_{\ell,k+1} , \forall \ell \in \mathbb{R}^+ , \forall k \in \mathbb{N}^* . \tag{A.3}$$

As a corollary, we obtain

$$j_{\ell,k} \leq j_{0,k+\ell} , \tag{A.4}$$

with strict inequality for  $\ell > 0$ .

It is also useful to have the half integer results (see in [1], p. 467)

$\ell =$	$\frac{1}{2}$	$\frac{3}{2}$	$\frac{5}{2}$	$\frac{7}{2}$	$\frac{9}{2}$	$\frac{11}{2}$	$\frac{13}{2}$
$k = 1$	$\pi$	4.49	5.76	6.99	8.18	9.36	10.51
2	$2\pi$	7.73	9.09	10.42	11.7		
3	$3\pi$	10.9	12.52	13.69	..		
4	$4\pi$	14.06					

(A.5)

## B Alternative approach in the regular case

Although not needed in this article, we recall some of the results of Statements of [16, 17]. The main theorem is the following

### Theorem B.1

Suppose that  $\Omega$  is regular and simply connected and that, for some regular closed set  $N$  satisfying in addition the “equal angle meeting condition”. Suppose that, for some  $\lambda \in \mathbb{R}$ , the associated family  $\mathcal{D}(N) = \{D_1, \dots, D_\mu\}$  is admissible and satisfies a Pair Compatibility Condition, which means that  $\lambda$  is an eigenvalue of  $H(D_{i,j})$  for which  $D_i$  and  $D_j$  are the two nodal domains of some corresponding eigenfunction. Then there is an eigenfunction  $u$  of  $H(\Omega)$  with corresponding eigenvalue  $\lambda$  such that the family of nodal domains of  $u$  is  $\mathcal{D}(N)$ .

### Remarks B.2

- If  $\Omega$  is not simply connected then the result does not hold in general. One should add a nonholonomy condition (see [17]). In the case of minimal partitions, we have seen that this condition is reduced to the bipartite condition.
- Note that the Pair Compatibility Condition is weaker than to assume that  $\lambda$  is the second eigenvalue of  $H(D_{i,j})$  for each pair of neighbors  $(D_i, D_j)$ .

As an application of this theorem, the authors obtain :

### Corollary B.3

Let  $\Omega$  be simply connected,  $k \in \mathbb{N}$  ( $k \geq 2$ ) and let  $\mathcal{D}^{min} = (D_i)_{i=1, \dots, k}$  be a minimal admissible strong regular<sup>6</sup> partition. Then there is an eigenfunction  $u$  of  $H(\Omega)$  associated with

$$\lambda = \max_i(\lambda(D_i)) ,$$

such that  $\mathcal{D}^{min}$  is the family of the  $k$  nodal domains of  $u$ .

### Proof

Let us apply Theorem B.1. We take as  $\lambda = \max_i(\lambda(D_i))$ .

---

<sup>6</sup>The notion of regularity was actually stronger there, but the Pair Compatibility Condition gives actually some regularity assumption on the boundaries.

The first point is that all the  $\lambda(D_i)$  should be equal. If not, one could by deformation of the  $D_i$ 's in a neighborhood of regular points of their boundary find a new partition  $\tilde{\mathcal{D}}$ , which would decrease  $\max_i(\lambda(D_i))$ .

The second point is to observe that considering two neighbors  $D_i$  and  $D_j$ , then  $\lambda$  should be the second eigenvalue of  $H(D_{i,j})$ . If it was not the case for some pair  $(i, j)$ , the two nodal domains of the second eigenfunction of  $H(D_{i,j})$  will give two new open sets  $D'_i$  and  $D'_j$  with  $\lambda(D'_i) = \lambda(D'_j)$ , in contradiction with the assumption of minimality and the first point of the proof.

Hence the Pair Compatibility Condition is satisfied.

**Remark B.4**

*As mentioned in the introduction, the case where  $k = 2$  corresponds to a rather well known characterization of the second eigenvalue of  $H(\Omega)$ . The admissibility condition is of course automatically satisfied in this case.*

## References

- [1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions*, Volume 55 of Applied Math Series. National Bureau of Standards, 1964.
- [2] G. Alessandrini. Critical points of solutions of elliptic equations in two variables. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 14(2):229–256 (1988).
- [3] G. Alessandrini. Nodal lines of eigenfunctions of the fixed membrane problem in general convex domains. *Comment. Math. Helv.*, 69(1):142–154 (1994).
- [4] A. Ancona, B. Helffer, and T. Hoffmann-Ostenhof. Nodal domain theorems à la Courant. *Documenta Mathematica* 9:283–299 (2004).
- [5] C. Bandle. Isoperimetric inequalities and applications. *Monographs and Studies in Mathematics*, 7, Pitman, 1980.
- [6] L. Bers. Local behaviour of solutions of general linear equations. *Commun. Pure Appl. Math.*, 8:473–496 (1955).
- [7] M. Conti, S. Terracini, and G. Verzini. An optimal partition problem related to nonlinear eigenvalues. *Journal of Functional Analysis* 198:160–196 (2003).
- [8] M. Conti, S. Terracini, and G. Verzini. A variational problem for the spatial segregation of reaction-diffusion systems. *Indiana Univ. Math. J.* 54(3):779–815 (2005).
- [9] M. Conti, S. Terracini, and G. Verzini. On a class of optimal partition problems related to the Fucik spectrum and to the monotonicity formula. *Calc. Var.* 22:45–72 (2005).
- [10] M. Dauge and B. Helffer. Eigenvalue variation II. Multidimensional problems. *Journal of differential equations* 104(2):263–297 (1993).
- [11] R. Diestel. Graph Theory, (second edition). *Graduate Texts in Mathematics*, 173, Springer 2000.
- [12] D. Gilbarg and N.S. Trudinger. Elliptic Partial Differential Equations of Second Order. *Springer* 1983.
- [13] M.J. Greenberg and J.R. Harper. *Algebraic topology*, volume 58 of *Mathematics Lecture Note Series*. Benjamin/Cummings Publishing Co. Inc. Advanced Book Program, Reading, Mass., 1981. A first course.
- [14] R. Hardt, M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and N. Nadirashvili. Critical sets of solutions to elliptic equations. *J. Differential Geom.* 51:359–373 (1999).
- [15] P. Hartman and A. Wintner. On the local behavior of solutions of non-parabolic partial differential equations. *Amer. J. Math.*, 75:449–476 (1953).
- [16] B. Helffer and T. Hoffmann-Ostenhof. Converse spectral problems for nodal domains. Extended version. Preprint Sept. 2005, mp\_arc 05343.
- [17] B. Helffer and T. Hoffmann-Ostenhof. Converse spectral problems for nodal domains. Preprint April 2006. To appear in *Moscow Mathematical Journal*.
- [18] B. Helffer and T. Hoffmann-Ostenhof. On nodal patterns and spectral optimal partitions. Unpublished Notes, December 2005.
- [19] M. Hoffmann-Ostenhof and T. Hoffmann-Ostenhof. Local properties solutions of Schrödinger equations *Comm. Partial Differential Equations* 17:491–522 (1992).

- [20] M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, and N. Nadirashvili. Interior Hölder estimates for solutions of Schrödinger equations and the regularity of nodal sets. *Comm. Partial Differential Equations* 20:1241–1243 (1995).
- [21] T. Hoffmann-Ostenhof, P. Michor, and N. Nadirashvili. Bounds on the multiplicity of eigenvalues for fixed membranes. *GAFSA*, 9:1169–1188 (1999).
- [22] T. Kato. Perturbation Theory for Linear Operators. Second edition. *Springer* 1977.
- [23] A. Pleijel. Remarks on Courant’s nodal theorem. *Comm. Pure. Appl. Math.*, 9: 543–550, 1956.
- [24] C. Pommerenke. Boundary behavior of conformal maps. Grundlehren der Mathematischen Wissenschaften 295, *Springer*, 1992.
- [25] M. Reed and B. Simon. Methods of modern mathematical physics IV: Analysis of operators. *Academic Press*, 1978.
- [26] B. Simon. Lower semi-continuity of positive quadratic forms. *Proc. Roy. Soc. Edin.* 29:267–273 (1977).
- [27] B. Simon. A canonical decomposition for quadratic forms with applications to monotone convergence theorems. *J. of Funct. Analysis* 28:377–385 (1978).
- [28] P. Stollmann. A convergence theorem for Dirichlet forms with applications to boundary value problems with varying domains. *Math. Z.*, 219: 275–287 (1995).
- [29] G.N. Watson. A treatise on the Theory of Bessel Functions. *Cambridge University Press* 1966.
- [30] J. Weidmann. Continuity of the eigenvalues of self-adjoint operators with respect to the strong operator topology. *Integral Equations Operator Theory* 3:138–142 (1980).
- [31] J. Weidmann. Monotone continuity of the spectral resolution and the eigenvalues. *Proc. R. Soc. Edimb. Sect. A*85:131–136 (1980).
- [32] J. Weidmann. Stetige Abhängigkeit der Eigenwerte und Eigenfunktionen elliptischer Differentialoperatoren vom Gebiet. *Math. Scand.* 54: 51–69 (1984).

B. HELFFER: DÉPARTEMENT DE MATHÉMATIQUES, BAT. 425, UNIVERSITÉ PARIS-SUD, 91 405 ORSAY CEDEX, FRANCE.

EMAIL: BERNARD.HELFFER@MATH.U-PSUD.FR

T. HOFFMANN-OSTENHOF: INSTITUT FÜR THEORETISCHE CHEMIE, UNIVERSITÄT WIEN, WÄHRINGER STRASSE 17, A-1090 WIEN, AUSTRIA AND INTERNATIONAL ERWIN SCHRÖDINGER INSTITUTE FOR MATHEMATICAL PHYSICS, BOLTZMANNGASSE 9, A-1090 WIEN, AUSTRIA.

EMAIL:THOFFMAN@ESI.AC.AT

S. TERRACINI: UNIVERSITÀ DI MILANO BICOCCA, VIA COZZI, 53 20125 MILANO (ITALY).

EMAIL: SUSANNA.TERRACINI@UNIMIB.IT